

BOX PATENT

Attorney Docket No. 22762XY4-S
IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of

Douglas T. ROSS et al.

Serial No. NOT YET ASSIGNED

Filed: August 14, 1998

For: CARBOXYLIC ACIDS AND ISOTERES OF N-HETEROCYCLIC COMPOUNDS FOR
VISION AND MEMORY DISORDERS

TRANSMITTAL LETTER

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

Submitted herewith for filing in the U.S. Patent and Trademark
Office is the following:

- (1) Transmittal Letter
- (2) PTO Form-1082
- (3) 122 page Application consisting of:
 - 102 pages of Textual Specification
 - 10 pages of 22 Claims
 - 1 Abstract of the Disclosure
 - 9 sheets of drawings
- (4) Executed Inventor's Declaration
- (5) Assignment with Recordation sheet
- (6) Check No. 10149 \$ 834.00 for Government Filing Fee
- (7) Check No. 10150 \$ 40.00 for Recordation of Assignment

Please charge any required fee, or credit any overpayment, in
connection with this matter to deposit Account No. 14-0112.

Respectfully submitted,


NATH & ASSOCIATES

By: _____

Gary M. Nath

Registration No. 26,965

Date: August 14 , 1998
NATH & ASSOCIATES
1835 K Street N.W., Suite 750
Washington, D.C. 20006-1203
GMN/ph



NATH & ASSOCIATES

Attorneys at Law

1835 K Street, N.W.

Suite 750

Washington, D.C. 20006-1203

TELEPHONE (202) 775-8383

FACSIMILE (202) 775-8396

(202) 822-9409

Patent, Trademark and Copyright Causes,
Unfair Competition, Trade Secrets,
Licensing and Antitrust

E-Mail: NATHINTELPROP@COMPUSERVE.COM

Gary M. Nath (NJ)

Irvin A. Lavine (DC)*

Karen Lee Orzechowski (DC, PA)

Jordan S. Weinstein (DC, NY, VA, PA, NJ)

Harold L. Novick (DC, MD)

Suet M. Chong (DC)

Todd L. Juneau (DC, IL)

Patricia M. Drost (DC, VA)

Gregory B. Kang (NJ)

Leigh A. Penfield (MD)

Michelle L. Hartland (VA)

Scott F. Yarnell (MD)

Paul Sacher (MD)

Lee C. Heiman (CA)

Robert G. Lev (DC)*

Donald M. Sandler (MD)

Alvin E. Tanenholtz**

*Of Counsel

**Registered Patent Agent

FORM PTO-1082

TRANSMITTAL FOR NEW U.S. PATENT APPLICATION

Assistant Commissioner ~~of~~ ~~for~~
~~Patents and Trademarks~~
Washington, D.C. 20231

Re: New U.S. Patent Application

For: CARBOXYLIC ACIDS AND ISOSTERES
OF N-HETEROCYCLIC COMPOUNDS FOR
VISION AND MEMORY DISORDERS

Inventor(s): Douglas T. ROSS et al.

Attorney Docket: 22762XY4-S

Sir:

Attached hereto is the application identified above, including:

122 Pages Application Consisting of:
102 Pages of Textual Specification
10 Pages of 22 Claims
1 Abstract of the Disclosure
2 Sheets of drawings.

☒ Executed Inventor's Declaration

☒ Assignment and cover sheet


The Government filing fee* is calculated as follows:

Total Claims	<u>22</u> - 20 = <u>2</u>	x \$22.00 =	<u>\$44.00</u>
Independent Claims	<u>2</u> - 3 = <u>0</u>	x \$80.00 =	<u>\$0.00</u>
Base Fee			<u>\$790.00</u>
Multiple Dependent Claim Fee (\$250.00)			<u>\$ 0.00</u>
Subtotal			<u>\$834.00</u>
TOTAL FILING FEE*			<u>\$834.00</u>

- ☐ *Reduced by one-half, as applicant(s) is/are a "small entity". A Declaration Claiming Small Entity Status:
 - ☐ will be filed at a later date.
- ☐ Foreign priority is claimed under 35 U.S.C. § 119 from Patent Application(s) No. dated .
 - ☐ Priority document(s) will be submitted at a later date.
 - ☐ Priority document(s) is/are submitted herewith.
- ☐ Executed Declaration(s) will be submitted at a later date pursuant to 37 CFR § 1.41 and § 1.53, with an appropriate surcharge under 37 CFR § 1.16(e).
- ☐ Formal drawing(s) is/are attached.
- ☒ Formal drawing(s) will be submitted at a later date.
- ☒ Assignment document(s) is/are submitted herewith; the recordation fee of \$40.00 per document is enclosed herewith.
- ☐ A Verified Translation will be submitted at a later date.
- ☐ No payment is enclosed at this time. Full payment will be made when the executed Declaration is submitted.
- ☒ Submitted herewith is a check in the amount of \$834.00. The Commissioner is hereby authorized to charge any deficiency or credit any excess to Deposit Account No. 14-0112.

Respectfully submitted,

NATH & ASSOCIATES



Gary M. Nath
Registration No. 26,965

Date: August 14, 1998

NATH & ASSOCIATES

1835 K Street, N.W., Suite 750

Washington, D.C. 20006-1203

(202)-775-8383 GMN/ph (1082)

Title:

CARBOXYLIC ACIDS AND ISOSTERES OF
N-HETEROCYCLIC COMPOUNDS
FOR VISION AND MEMORY DISORDERS

Inventors: Douglas T. Ross
Hansjörg Sauer
Gregory S. Hamilton
Joseph P. Steiner

NATH & ASSOCIATES

1835 K Street, N.W., Suite 750

Washington, D.C. 20006-1203

202-775-8383

BACKGROUND OF THE INVENTION

5

1. Field of Invention

10 This invention relates to pharmaceutical compositions and methods for treating vision loss, preventing vision degeneration, and promoting vision regeneration ("neopsis") using low molecular weight, small molecule derivatives.

2. Description of Related Art

15 The visual system is composed of the eyes, ocular adnexa and the visual pathways. Dysfunction of the visual system may lead to permanent or temporary visual impairment, i.e. a deviation from normal in one or more functions of the eye. Visual impairment manifests itself in various ways and includes a broad
20 range of visual dysfunctions and disturbances. Without limitation, these dysfunctions and disturbances include partial or total loss of vision, the need for correction of visual acuity for objects near and far, loss of visual field, impaired ocular
25 motility without diplopia (double vision), impaired or skewed color perception, limited adaptation to light and dark, diminished accommodation, metamorphopsic

distortion, impaired binocular vision, paresis of accommodation, iridoplegia, entropion, ectropion, epiphora, lagophthalmos, and scarring. See

Physicians' Desk Reference (PDR) for Ophthalmology,

5 16th Edition, 6:47 (1988). The visual system may be adversely affected by various ophthalmologic disorders, diseases, injuries, and complications, including, without limitation, genetic disorders; [non-genetic disorders;] disorders associated with
10 aging or degenerative diseases; disorders correlating to physical injury to the eye, head, or other parts of the body resulting from external forces; disorders resulting from environmental factors; disorders resulting from a broad range of diseases; and
15 combinations of any of the above.

The visual system is a complex system composed of numerous components. Visual impairment can involve the entire visual system, any one component, or any combination of components, depending upon the precise
20 nature of the circumstances. The eye is composed of a lens, which is suspended in the zonules of Zinn and is focused by the ciliary body. The ciliary body also secretes aqueous humor, which fills the posterior chamber, passes through the pupil into the anterior
25 chamber, then drains primarily via the canal of Schlemm. The iris regulates the quantity of light

entering the eye by adjusting the size of its central opening, the pupil. A visual image is focused onto the retina, the fovea centralis being the retinal area of sharpest visual acuity. The conjunctiva is the mucus membrane which lines the eyelids and the eyeball, and ends abruptly at the limbus conjunctivae, the edge of the conjunctiva overlapping the cornea. The cornea is the clear, transparent anterior portion of the fibrous coat of the eye; it is important in light refraction and is covered with an epithelium that differs in many respects from the conjunctival epithelium.

The retina is the innermost, light sensitive portion of the eye, containing two types of photoreceptors, cones, which are responsible for color vision in brighter light, and rods, which are essential for vision in dim light but do not perceive colors. After light passes through the cornea, lens system, and the vitreous humor, it enters the retina from the inside; that is, it passes through the ganglion cells and nerve fibers, the inner and outer plexiform layers, the inner and outer nuclear layers, and the internal and external limiting membranes before it finally reaches the layer of photoreceptors located near the outside of the retina, just inside the outermost pigment epithelium layer. The cells of

the pigment epithelium layer act as an anatomical barrier to liquids and substances located outside of the eye, forming the "blood-retina" barrier, and provide nourishment, oxygen, a source of functionally useful substances like vitamin A, and phagocytosis of decomposition products to photoreceptor cells. There is no anatomical connection between the pigment epithelium and the photoreceptor layer, permitting separation of the layers in some pathological situations.

When rods or cones are excited by light, signals are transmitted through successive neurons in the retina itself, into the optic nerve fibers, and ultimately to the cerebral cortex. Both rods and cones contain molecules that decompose on exposure to light and, in the process, excite the nerve fibers leading from the eye. The molecule in rods is rhodopsin. The three light-sensitive molecules in cones, collectively called iodopsin, have compositions only slightly different from that of rhodopsin and are maximally excited by red, blue, or green light, respectively.

Neither rods nor cones generate action potentials. Rather, the light-induced membrane hyperpolarization generated in the outer, photosensitive segment of a rod or cone cell is

transmitted from the outer segment through the inner
segment to the synaptic body by direct conduction of
the electrical voltage itself, a process called
electrotonic conduction. At the synaptic body, the
5 membrane potential controls the release of an unknown
transmitter molecule. In low light, rod and cone cell
membranes are depolarized and the rate of transmitter
release is greatest. Light-induced hyperpolarization
causes a marked decrease in the release of transmitter
10 molecules.

The transmitters released by rod and cone cells
induce signals in the bipolar neurons and horizontal
cells. The signals in both these cells are also
transmitted by electrotonic conduction and not by
15 action potential.

The rod bipolar neurons connect with as many as
50 rod cells, while the dwarf and diffuse bipolar
cells connect with one or several cone cells. A
depolarizing bipolar cell is stimulated when its
20 connecting rods or cones are exposed to light. The
release of transmitter molecules inhibits the
depolarizing bipolar cell. Therefore, in the dark,
when the rods and cones are secreting large quantities
of transmitter molecules, the depolarizing bipolar
25 cells are inhibited. In the light, the decrease in
release of transmitter molecules from the rods and

cones reduces the inhibition of the bipolar cell, allowing it to become excited. In this manner, both positive and negative signals can be transmitted through different bipolar cells from the rods and
5 cones to the amacrine and ganglion cells.

As their name suggests, horizontal cells project horizontally in the retina, where they may synapse with rods, cones, other horizontal cells, or a combination of cells types. The function of
10 horizontal cells is unclear, although some mechanism in the convergence of photoreceptor signaling has been postulated.

All types of bipolar cells connect with ganglion cells, which are of two primary types. A-type
15 ganglion cells predominately connect with rod bipolar cells, while B-type ganglion cells predominately connect with dwarf and diffuse bipolar cells. It appears that A-type ganglion cells are sensitive to contrast, light intensity, and perception of movement,
20 while B-type ganglion cells appear more concerned with color vision and visual acuity.

Like horizontal cells, the Amacrine cells horizontally synapse with several to many other cells, in this case bipolar cells, ganglion cells, and other
25 Amacrine cells. The function of Amacrine cells is also unclear.

The axons of ganglion cells carry signals into the nerve fiber layer of the eye, where the axons converge into fibers which further converge at the optic disc, where they exit the eye as the optic nerve. The ganglion cells transmit their signals through the optic nerve fibers to the brain in the form of action potentials. These cells, even when unstimulated, transmit continuous nerve impulses at an average, baseline rate of about 5 per second. The visual signal is superimposed onto this baseline level of ganglion cell stimulation. It can be either an excitatory signal, with the number of impulses increasing above the baseline rate, or an inhibitory signal, with the number of nerve impulses decreasing below the baseline rate.

As part of the central nervous system, the eye is in some ways an extension of the brain; as such, it has a limited capacity for regeneration. This limited regeneration capacity further complicates the challenging task of improving vision, resolving dysfunction of the visual system, and/or treating or preventing ophthalmologic disorders. Many disorders of the eye, such as retinal photic injury, retinal ischemia-induced eye injury, age-related macular degeneration, free radical-induced eye diseases, as well as numerous other disorders, are considered to be

entirely untreatable. Other ophthalmologic disorders, e.g., disorders causing permanent visual impairment, are corrected only by the use of ophthalmic devices and/or surgery, with varying degrees of success.

5 The immunosuppressant drugs FK506, rapamycin, and cyclosporin are well known as potent T-cell specific immunosuppressants, and are effective against autoimmunity, transplant or graft rejection, inflammation, allergic responses, other autoimmune or
10 immune-mediated diseases, and infectious diseases. It has been disclosed that application of Cyclosporin, FK-506, Rapamycin, Buspirone, Spiperone, and/or their derivatives are effective in treating some ophthalmologic disorders of these types. Several
15 ophthalmologic disorders or vision problems are known to be associated with autoimmune and immunologically-mediated activities; hence, immunomodulatory compounds are expected to demonstrate efficacy for treating those types of ophthalmologic disorders or vision
20 problems.

 The effects of FK506, Rapamycin, and related agents in the treatment of ophthalmologic diseases are disclosed in several U.S. patents (Goulet et al., U.S. Patent No. 5,532,248; Mochizuki et al., U.S. Patent
25 No. 5,514,686; Luly et al., U.S. Patent No. 5,457,111; Russo et al., U.S. Patent No. 5,441,937; Kulkarni,

U.S. Patent No. 5,387,589; Asakura et al., U.S. Patent No. 5,368,865; Goulet et al., U.S. Patent No. 5,258,389; Armistead et al., U.S. Patent No. 5,192,773; Goulet et al., U.S. Patent No. 5,189,042; and Fehr, U.S. Patent No. 5,011,844). These patents claim FK506 or Rapamycin related compounds and disclose the known use of FK506 or Rapamycin related compounds in the treatment of ophthalmologic disorders in association with the known immunosuppressive effects of FK506 and Rapamycin. The compounds disclosed in these patents are relatively large. Further, the cited patents relate to immunomodulatory compounds limited to treating autoimmunity or related diseases, or immunologically-mediated diseases, for which the efficacy of FK506 and Rapamycin is well known.

Other U.S. patents disclose the use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds for use in the treatment of ophthalmologic diseases (Sharpe et al., U.S. Patent No. 5,703,088; Sharpe et al., U.S. Patent No. 5,693,645; Sullivan, U.S. Patent No. 5,688,765; Sullivan, U.S. Patent No. 5,620,921; Sharpe et al., U.S. Patent No. 5,574,041; Eberle, U.S. Patent No. 5,284,826; Sharpe et al., U.S. Patent No. 5,244,902; Chiou et al., U.S. Patent Nos. 5,198,454 and

5,194,434; and Kaswan, U.S. Patent No. 4,839,342).

These patents also relate to compounds useful for treating autoimmune diseases and cite the known use of cyclosporin, Spiperone, Buspirone, their derivatives, and other immunosuppressive compounds in treating ocular inflammation and other immunologically-mediated ophthalmologic diseases.

The immunosuppressive compounds disclosed in the prior art suppress the immune system, by definition, and also exhibit other toxic side effects. Accordingly, there is a need for non-immunosuppressant, small molecule compounds, and compositions and methods for use of such compounds, that are useful in improving vision; preventing, treating, and/or repairing visual impairment or dysfunction of the visual system; and preventing, treating, and/or resolving ophthalmologic disorders.

There are also a number of patents on non-immunosuppressive compounds disclosing methods of use for permitting or promoting wound healing (whether from injury or surgery); controlling intraocular pressure (often resulting from glaucoma); controlling neurodegenerative eye disorders, including damage or injury to retinal neurons, damage or injury to retinal ganglion cells, and macular degeneration; stimulating neurite outgrowth; preventing or reducing oxidative

damage caused by free radicals; and treating impaired oxygen and nutrient supply, as well as impaired waste product removal, resulting from low blood flow. These non-immunosuppressive substances fall into one of two
5 general categories: naturally occurring molecules, such as proteins, glycoproteins, peptides, hormones, and growth factors; and synthetic molecules.

Within the group of naturally occurring non-immunosuppressive molecules, several hormones, growth
10 factors, and signaling molecules have been patented for use as supplements to naturally occurring quantities of such molecules, as well as for targeting of specific cells where the particular molecule does not naturally occur in a mature individual. These
15 patents generally claim methods of use for reducing or preventing the symptoms of ocular disease, or arresting or reversing vision loss.

Specifically, Louis et al., U.S. Patent Nos. 5,736,516 and 5,641,749, disclose the use of a glial
20 cell line derived neurotrophic factor (GDNF) to stop or reverse the degeneration of retinal neurons (i.e. photoreceptors) and retinal ganglion cells caused by glaucoma, or other degenerative or traumatic retinal diseases or injuries. O'Brien, et al., U.S. Patent
25 Nos. 5,714,459 and 5,700,909, disclose the use of a glycoprotein, Saposin, and its derivatives for

stimulating neurite outgrowth and increasing myelination. To stop or reverse degeneration of retinal neurons, LaVail et al., U.S. Patent No. 5,667,968, discloses the use of a variety of neurotrophic proteins, including brain-derived neurotrophic factor, ciliary neurotrophic factor, neurotrophin-3 or neurotrophin-4, acidic or basic fibroblast growth factors, interleukin, tumor necrosis factor- α , insulin-like growth factor-2 and other growth factors. Wong et al., U.S. Patent No. 5,632,984, discloses the use of interferons, especially interferon α -2a, for treating the symptoms of macular degeneration by reducing hemorrhage and limiting neovascularization. Finally, Wallace et al., U.S. Patent No. 5,441,937, discloses the use of a lung-derived neurotrophic factor (NTF) to maintain the functionality of ciliary ganglion and parasympathetic neuron cells.

A key characteristic of factors derived from specific cell lines is their localization to specific cell lines or tissues; systemic treatment with these molecules would run a substantial risk of unintended, and potentially dangerous, effects in cell lines where the genes encoding these molecules are inactive. Similarly, hormones and growth factors often activate a large number of genes in many cell lines; again,

non-localized application of these molecules would run a substantial risk of provoking an inappropriate, and potentially dangerous, response.

Within the category of synthetic molecules, most of the patented compounds are immunosuppressive and disclose uses in treating inflammatory, autoimmune, and allergic responses, as discussed above. A few others are non-immunosuppressive and claim the ability to treat cellular degeneration, and in some cases promote cellular regeneration, most often in the context of their antioxidant properties.

Specifically, Tso et al., U.S. Patent No. 5,527,533, discloses the use of astaxanthin, a carotenoid antioxidant, for preventing or reducing photoreceptor damage resulting from the presence of free radicals. Similarly, Babcock et al., U.S. Patent No. 5,252,319, discloses the use of antioxidant aminosteroids for treating eye disease and injury, by increasing resistance to oxidative damage. Freeman, U.S. Patent No. 5,468,752, discloses the use of the antiviral phosphonylmethoxyalkylcytosines to reduce abnormally increased intraocular pressure.

Hamilton and Steiner disclose in U.S. Patent No. 5,614,547 novel pyrrolidine carboxylate compounds which bind to the immunophilin FKBP12 and stimulate nerve growth, but which lack immunosuppressive


effects. Unexpectedly, it has been discovered that these non-immunosuppressant compounds promote improvements in vision and resolve ophthalmologic disorders. Yet their novel small molecule structure and non-immunosuppressive properties differentiate them from FK506 and related immunosuppressive compounds found in the prior art.

Further, these compounds may be differentiated from the non-immunosuppressive compounds used to treat vision disorders by their novel small molecule structure and their lack of general, systemic effects. Naturally occurring hormones, growth factors, cytokines, and signaling molecules are generally multifunctional and activate many genes in diverse cell lines. The present compounds do not, thus avoiding the unexpected, and potentially dangerous, side effects of systemic use. Similarly, the present compounds also avoid the potential unexpected side effects of introducing cell line-specific molecules into other cell lines were they do not naturally occur.

SUMMARY OF THE INVENTION

The present invention relates to the surprising discovery that N-heterocyclic compounds containing a carboxylic acid or isostere thereof may be useful for treating a vision disorder, improving vision, treating memory impairment or enhancing memory performance in an animal. Accordingly, novel compositions and methods of using N-heterocyclic ring compounds containing an acidic moiety or an isostere thereof attached to the 2-carbon of the N-heterocyclic ring are provided. A preferred feature of the compounds of the present invention is that they do not exert any significant immunosuppressive activity.

Preferred embodiments of this invention include methods and compositions containing a compound having the formula (I):



I

where

n is 1-3;

X is either O or S;


R₁ is selected from the group consisting of C₁-C₉,
straight or branched chain alkyl, C₂-C₉, straight or
5 branched chain alkenyl, aryl, heteroaryl, carbocycle, or
heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain
alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl; and

R₂ is a carboxylic acid or a carboxylic acid isostere;
10 or a pharmaceutically acceptable salt, ester, or solvate
thereof;

Preferred embodiments of this invention are where R₂
is a carbocycle or heterocycle containing any combination
of CH₂, O, S, or N in any chemically stable oxidation
15 state, where any of the atoms of said ring structure are
optionally substituted in one or more positions with R³.

Especially preferred embodiments of this invention
are where R₂ is selected from the group below:



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 ,

where

- 5 R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl or
10 alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and CO_2R^4 where R^4 is hydrogen or C_1 - C_6 straight or branched chain alkyl or alkenyl.

- Another preferred embodiment of this invention is where R_2 is selected from the group consisting of -COOH,
15 - SO_3H , - SO_2HNR^3 , - $PO_2(R^3)_2$, -CN, - $PO_3(R^3)_2$, - OR^3 , - SR^3 , - $NHCO R^3$, - $N(R^3)_2$, - $CON(R^3)_2$, - $CONH(O)R^3$, - $CONHNHSO_2R^3$, - $COHNSO_2R^3$, and - $CONR^3CN$.

- Preferred compounds of this invention are: (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-hydroxymethyl
20 pyrrolidine; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinetetrazole; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile; and (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-aminocarbonyl
piperidine, and compounds 1-136 of Tables I, II, and III.

Brief Description of the Drawings

5 Figure 1 A, B and C show that GPI 1046 protects retinal ganglion cells against degeneration following retinal ischemia.

10 Figure 2 shows that GPI 1046 prevents degeneration of optic nerve axons and myelin following retinal ischemia.

15 Figure 3 shows that GPI 1046 provides moderate protection against retinal ganglion cell death after optic nerve transection.

20 Figure 4 shows that GPI 1046 treatment duration significantly affects the process of optic nerve axonal degeneration after transection.

Figure 5 shows that GPI 1046 treatment produces a greater effect on optic nerve axons than ganglion cell bodies.

25 Figure 6 shows that GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in the proximal stump.

Figure 7 shows that FKBP-12 immunohistochemistry labels oligodendroglia (large dark cells with fibrous processes), the cells which produce myelin, located
5 between the fascicles of optic nerve fibers, and also some optic nerve axons.

Figure 8 shows GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in
10 the distal stump.

Figure 9 shows that 28 day treatment with GPI 1046 treatment beginning 8 weeks after onset of streptozotocin induced diabetes decreases the extent of
15 neovascularization in the inner and outer retina and protects neurons in the inner nuclear layer (INL) and ganglion cell layer (GCL) from degeneration.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

"Eye" refers to the anatomical structure responsible for vision in humans and other animals, and encompasses the following anatomical structures, without limitation: lens, vitreous body, ciliary body, posterior chamber, anterior chamber, pupil, cornea, iris, canal of Schlemm, zonules of Zinn, limbus, conjunctiva, choroid, retina, central vessels of the retina, optic nerve, fovea centralis, macula lutea, and sclera.

"Alkyl" means a branched or unbranched saturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₁-C₆ straight or branched alkyl hydrocarbon chain contains 1 to 6 carbon atoms, and includes but is not limited to substituents such as methyl, ethyl, propyl, iso-propyl, butyl, iso-butyl, tert-butyl, n-pentyl, n-hexyl, and the like. It is also contemplated as within the scope of the present invention that "alkyl" may also refer to a hydrocarbon chain wherein any of the carbon atoms of said alkyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of n-pentyl can be replaced with O to form propyloxymethyl.

"Alkenyl" means a branched or unbranched unsaturated hydrocarbon chain comprising a designated number of carbon atoms. For example, C₂-C₆ straight or branched

alkenyl hydrocarbon chain contains 2 to 6 carbon atoms having at least one double bond, and includes but is not limited to substituents such as ethenyl, propenyl, isopropenyl, butenyl, iso-butenyl, tert-butenyl, n-pentenyl, n-hexenyl, and the like. It is also contemplated as within the scope of the present invention that "alkenyl" may also refer to an unsaturated hydrocarbon chain wherein any of the carbon atoms of said alkenyl are optionally replaced with O, NH, S, or SO₂. For example, carbon 2 of 4-pentene can be replaced with O to form (2-propene)oxymethyl.

"Alkoxy" means the group -OR wherein R is alkyl as herein defined. Preferably, R is a branched or unbranched saturated hydrocarbon chain containing 1 to 6 carbon atoms.

"Aryl, heteroaryl, carbocycle, or heterocycle" includes but is not limited to cyclic or fused cyclic ring moieties and includes a mono-, bi- or tricyclic, carbo- or heterocyclic ring, wherein the ring is either unsubstituted or substituted in one or more position(s) with hydroxy, carbonyl, amino, amido, cyano, isocyan, nitro, nitroso, nitrilo, isonitrilo, imino, azo, diazo, sulfonyl, sulfhydryl, sulfoxy, thio, thiocarbonyl, thiocyano, formanilido, thioformamido, sulfhydryl, halo, haloalkyl, trifluoromethyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, alkylamino,

aminoalkyl, thioalkyl, alkylthio, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, or CO₂R⁴ where R⁴ is hydrogen or C₁-C₆ straight or branched chain alkyl and carbocyclic and heterocyclic moieties; wherein the individual ring sizes are 5-8 members; wherein the heterocyclic ring contains 1-4 heteroatom(s) selected from the group consisting of O, N, or S; wherein aromatic or tertiary alkyl amines are optionally oxidized to a corresponding N-oxide.

Examples of useful alkyl groups include, without limitation, methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, n-pentyl, 2-methyl pentyl and the like.

Examples of useful carbocyclic and heterocyclic moieties include, without limitation, phenyl, benzyl, naphthyl, indenyl, azulenyl, fluorenyl, anthracenyl, indolyl, isoindolyl, indolinyl, benzofuranyl, benzothiophenyl, indazolyl, benzimidazolyl, benzthiazolyl, tetrahydrofuranyl, tetrahydropyranyl, pyridyl, pyrrolyl, pyrrolidinyl, pyridinyl, pyrimidinyl, purinyl, quinc₁inyl, isoquinolinyl, tetrahydroquinolinyl, quinoliziny₁, furyl, thiophenyl, imidazolyl, oxazolyl, benzoxazolyl, thiazolyl, isoxazolyl, isotriazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, trithianyl, indoliziny₁, pyrazolyl, pyrazolinyl, pyrazolidinyl,

thienyl, tetrahydroisoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, naphthyridinyl, pteridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl, phenoxazinyl, and adamantyl.

5 "Halo" means at least one fluoro, chloro, bromo, or iodo moiety.

10 The term "pharmaceutically acceptable salt, ester, or solvate" refers to salt, ester, or solvates of the subject compounds which possess the desired pharmacological activity and which are neither biologically nor otherwise undesirable. The salt, ester, or solvates can be formed with inorganic or organic acids such as acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucoheptanoate, gluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, naphthylate, 2-naphthalenesulfonate, nicotinate, oxalate, sulfate, thiocyanate, tosylate and undecanoate. Base salt, ester, or solvates include ammonium salts, alkali metal salts such as lithium, sodium and potassium salts, alkaline earth metal salts such as calcium and magnesium salts, salt with organic bases such as dicyclohexylamine salts,

15
20
25

N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and so forth. Also, the basic nitrogen-containing groups can be quarternized with such agents as: 1) lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; 2) dialkyl sulfates like dimethyl, diethyl, dibutyl and diamyl sulfates; 3) long chain alkyls such as decyl, lauryl, myristyl and stearyl substituted with one or more halide such as chloride, bromide and iodide; and 4) aralkyl halides like benzyl and phenethyl bromide and others.

The compounds of this invention may possess at least one asymmetric center and thus can be produced as mixtures of stereoisomers or as individual enantiomers or diastereomers. The individual stereoisomers may be obtained by using an optically active starting material, by resolving a racemic or non-racemic mixture of an intermediate at some appropriate stage of the synthesis, or by resolution of the compound of formula (I). It is understood that the individual stereoisomers as well as mixtures (racemic and non-racemic) of stereoisomers are encompassed by the scope of the present invention. The S-stereoisomer at atom 1 of formula I is a most preferred embodiment of the invention.

"Stereoisomers" are isomers that differ only in the way the atoms are arranged in space.

"Isomers" are different compounds that have the same molecular formula and includes cyclic isomers such as (iso)indole and other isomeric forms of cyclic moieties.


"Enantiomers" are a pair of stereoisomers that are non-superimposable mirror images of each other.

"Diastereoisomers" are stereoisomers which are not mirror images of each other.

"Racemic mixture" means a mixture containing equal parts of individual enantiomers. "Non-racemic mixture" is a mixture containing unequal parts of individual enantiomers or stereoisomers.

"Isosteres" are different compounds that have different molecular formulae but exhibit the same or similar properties. For example, tetrazole is an isostere of carboxylic acid because it mimics the properties of carboxylic acid even though they both have very different molecular formulae. Tetrazole is one of many possible isosteric replacements for carboxylic acid. Other carboxylic acid isosteres contemplated by the present invention include $-\text{COOH}$, $-\text{SO}_3\text{H}$, $-\text{SO}_2\text{HNR}^3$, $-\text{PO}_2(\text{R}^3)_2$, $-\text{CN}$, $-\text{PO}_3(\text{R}^3)_2$, $-\text{OR}^3$, $-\text{SR}^3$, $-\text{NHCOR}^3$, $-\text{N}(\text{R}^3)_2$, $-\text{CON}(\text{R}^3)_2$, $-\text{CONH}(\text{O})\text{R}^3$, $-\text{CONHNHSO}_2\text{R}^3$, $-\text{COHNSO}_2\text{R}^3$, and $-\text{CONR}^3\text{CN}$. In addition, carboxylic acid isosteres can include 5-7 membered carbocycles or heterocycles containing any combination of CH_2 , O, S, or N in any chemically stable oxidation state, where any of the atoms of said ring

structure are optionally substituted in one or more positions. The following structures are non-limiting examples of preferred carbocyclic and heterocyclic isosteres contemplated by this invention.



5

where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 . The present invention contemplates that when chemical substituents are added to a carboxylic isostere then the inventive compound retains the properties of a carboxylic isostere. The present invention contemplates that when a carboxylic isostere is optionally substituted with one or more moieties selected from R^3 , then the substitution cannot eliminate the carboxylic acid isosteric properties of the inventive compound. The present invention contemplates that the placement of one or more R^1 substituents upon a carbocyclic or heterocyclic carboxylic acid isostere shall not be permitted at one or more atom(s) which maintain(s) or is/are integral to the carboxylic acid isosteric properties of the inventive compound, if such substituent(s) would destroy the carboxylic acid isosteric properties of the inventive compound.

Other carboxylic acid isosteres not specifically exemplified or described in this specification are also contemplated by the present invention.

The term "preventing neurodegeneration" as used herein includes the ability to inhibit or prevent neurodegeneration in patients newly diagnosed as having a neurodegenerative disease, or at risk of developing a new degenerative disease and for inhibiting or preventing further neurodegeneration in patients who are already suffering from or have symptoms of a neurodegenerative disease.

The term "treatment" as used herein covers any treatment of a disease and/or condition in an animal, particularly a human, and includes:

(i) preventing a disease and/or condition from occurring in a subject which may be predisposed to the disease and/or condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease and/or condition, i.e., arresting its development; or

(iii) relieving the disease and/or condition, i.e., causing regression of the disease and/or condition.

The system used in naming the compounds of the present invention is shown below, using a compound of formula I as an example.

A compound of the present invention, especially formula I, wherein n is 1, X is O, D is a bond, R₁ is 1,1-dimethylpropyl, and R₂ is -CN, is named (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile.

15 "Enhancing memory performance" refers to
improving or increasing the mental faculty by which to
register, retain or recall past experiences,
knowledge, ideas, sensations, thoughts or impressions.

20 "Memory impairment" refers to a diminished mental
registration, retention or recall of past experiences,
knowledge, ideas, sensations, thoughts or impressions.
Memory impairment may affect short and long-term
information retention, facility with spatial
relationships, memory (rehearsal) strategies, and
verbal retrieval and production. Common causes of

memory impairment are age, severe head trauma, brain anoxia or ischemia, alcoholic-nutritional diseases, and drug intoxications. Examples of memory impairment include, without limitation, benign forgetfulness, amnesia and any disorder in which memory deficiency is present, such as Korsakoff's amnesic psychosis, dementia and learning disorders.

"Neopsic factors" or "neopsics" refers to compounds useful in treating vision loss, preventing vision degeneration, or promoting vision regeneration.

"Neopsis" refers to the process of treating vision loss, preventing vision degeneration, or promoting vision regeneration.

"Ophthalmological" refers to anything about or concerning the eye, without limitation, and is used interchangeably with "ocular," "ophthalmic," "ophthalmologic," and other such terms, without limitation.

"Preventing vision degeneration" refers to the ability to prevent degeneration of vision in patients newly diagnosed as having a degenerative disease

affecting vision, or at risk of developing a new degenerative disease affecting vision, and for preventing further degeneration of vision in patients who are already suffering from or have symptoms of a
5 degenerative disease affecting vision.

"Promoting vision regeneration" refers to maintaining, improving, stimulating or accelerating recovery of, or revitalizing one or more components of the visual system in a manner which improves or
10 enhances vision, either in the presence or absence of any ophthalmologic disorder, disease, or injury.

"Treating" refers to:

(i) preventing a disease and/or condition from occurring in a subject which may be predisposed to the
15 disease and/or condition but has not yet been diagnosed as having it;

(ii) inhibiting the disease and/or condition, i.e., arresting its development; or

(iii) relieving the disease and/or condition,
20 i.e., causing regression of the disease and/or condition.

"Vision" refers to the ability of humans and other animals to process images, and is used interchangeably with "sight", "seeing", and other such
25 terms, without limitation.

"Vision disorder" refers to any disorder that affects or involves vision, including without

limitation visual impairment, orbital disorders, disorders of the lacrimal apparatus, disorders of the eyelids, disorders of the conjunctiva, disorders of the cornea, cataracts, disorders of the uveal tract, disorders of the retina, disorders of the optic nerve or visual pathways, free radical induced eye disorders and diseases, immunologically-mediated eye disorders and diseases, eye injuries, and symptoms and complications of eye disease, eye disorder, or eye injury.

"Visual impairment" refers to any dysfunction in vision including without limitation disturbances or diminution in vision (e.g., binocular, central, peripheral, scotopic), visual acuity for objects near and far, visual field, ocular motility, color perception, adaptation to light and dark, accommodation, refraction, and lacrimation. See Physician's Desk Reference (PDR) for Ophthalmology, 16th Edition, 6:47 (1988).

20

Methods of the Present Invention

The present invention relates to a method of treating a vision disorder, improving vision, treating memory impairment, or enhancing memory performance in an animal, which comprises administering to said animal an effective amount of a derivative.

The inventive methods are particularly useful for treating various eye disorders including but not limited to visual disorders, diseases, injuries, and complications, genetic disorders; disorders associated with aging or degenerative vision diseases; vision disorders correlating to physical injury to the eye, head, or other parts of the body resulting from external forces; vision disorders resulting from environmental factors; vision disorders resulting from a broad range of diseases; and combinations of any of the above.

In particular, the compositions and methods of the present invention are useful for improving vision, or correcting, treating, or preventing visual (ocular) impairment or dysfunction of the visual system, including permanent and temporary visual impairment, without limitation. The present invention is also useful in preventing and treating ophthalmologic diseases and disorders, treating damaged and injured eyes, and preventing and treating diseases, disorders, and injuries which result in vision deficiency, vision loss, or reduced capacity to see or process images, and the symptoms and complications resulting from same. The eye diseases and disorders which may be treated or prevented by the compositions and methods of the present invention are not limited with regard to the cause of said diseases or disorders.

Accordingly, said compositions and methods are applicable whether the disease or disorder is caused by genetic or environmental factors, as well as any other influences. The compositions and methods of the present invention are particularly useful for eye problems or vision loss or deficiency associated with all of the following, without limitation: aging, cellular or physiological degeneration, central nervous system or neurological disorder, vascular defects, muscular defects, and exposure to adverse environmental conditions or substances.

The compositions and methods of the present invention are particularly useful in correcting, treating, or improving visual impairment, without limitation. Visual impairment in varying degrees occurs in the presence of a deviation from normal in one or more functions of the eye, including (1) visual acuity for objects at distance and near; (2) visual fields; and (3) ocular motility without diplopia. See *Physicians' Desk Reference (PDR) for Ophthalmology, 16th Edition, 6:47 (1988)*. Vision is imperfect without the coordinated function of all three. *Id.*

Said compositions and methods of use are also useful in correcting, treating, or improving other ocular functions including, without limitation, color perception, adaptation to light and dark, accommodation, metamorphopsia, and binocular vision.

The compositions and methods of use are particularly useful in treating, correcting, or preventing ocular disturbances including, without limitation, paresis of accommodation, iridoplegia, entropion, ectropion, 5 epiphora, lagophthalmos, scarring, vitreous opacities, non-reactive pupil, light scattering disturbances of the cornea or other media, and permanent deformities of the orbit.

The compositions and methods of use of the 10 present invention are also highly useful in improving vision and treating vision loss. Vision loss ranging from slight loss to absolute loss may be treated or prevented using said compositions and methods of use. Vision may be improved by the treatment of eye 15 disorders, diseases, and injuries using the compositions and methods of the invention. However, improvements in vision using the compositions and methods of use are not so limited, and may occur in the absence of any such disorder, disease, or injury.

20 The compositions and methods of the present invention are also useful in the treatment or prevention of the following non-limiting exemplary diseases and disorders, and symptoms and complications resulting therefrom.

25 Vision disorders include but are not limited to the following:

visual impairment, such as diminished visual

acuity for objects near and far, visual fields, and ocular motility;

orbital disorders, such as orbital cellulitis, periorbital cellulitis, cavernous sinus thrombosis, and exophthalmos (proptosis);

disorders of the lacrimal apparatus, such as dacryostenosis, congenital dacryostenosis, and dacryocystitis (acute or chronic);

disorders of the eyelids, such as lid edema, blepharitis, ptosis, Bell's palsy, blepharospasm, hordeolum (stye), external hordeolum, internal hordeolum (meibomian stye), chalazion, entropion (inversion of the eyelid), ectropion (eversion of the eyelid), tumors (benign and malignant), xanthelasma, basil cell carcinoma, squamous cell carcinoma, meibomian gland carcinoma, and melanoma;

disorders of the conjunctiva, such as pinguecula, pterygium, and other neoplasms, acute conjunctivitis, chronic conjunctivitis, adult gonococcal conjunctivitis, neonatal conjunctivitis, trachoma (granular conjunctivitis or Egyptian ophthalmia), inclusion conjunctivitis (inclusion blenorrhea or swimming pool conjunctivitis), neonatal inclusion conjunctivitis, adult inclusion conjunctivitis, vernal keratoconjunctivitis, keratoconjunctivitis sicca (keratitis sicca or dry eye syndrome), episcleritis, scleritis, cicatricial pemphigoid (ocular cicatricial

pemphigoid or benign mucous membrane pemphigoid), and subconjunctival hemorrhage;

disorders of the cornea, such as superficial punctate keratitis, corneal ulcer, indolent ulcer, recurrent corneal erosion, corneal epithelial basement membrane dystrophy, corneal endothelial cell dystrophy, herpes simplex keratitis (herpes simplex keratoconjunctivitis), dendritic keratitis, disciform keratitis, ophthalmic herpes zoster, phlyctenular keratoconjunctivitis (phlyctenular or eczematous conjunctivitis), interstitial keratitis (parenchymatous keratitis), peripheral ulcerative keratitis (marginal keratolysis or peripheral rheumatoid ulceration), keratomalacia (xerotic keratitis), xerophthalmia, keratoconus, bullous keratopathy;

cataracts, including developmental or congenital cataracts, juvenile or adult cataracts, nuclear cataract, posterior subcapsular cataracts;

disorders of the uveal tract, such as uveitis (inflammation of the uveal tract or retina), anterior uveitis, intermediate uveitis, posterior uveitis, iritis, cyclitis, choroiditis, ankylosing spondylitis, Reiter's syndrome, pars planitis, toxoplasmosis, cytomegalovirus (CMV), acute retinal necrosis, toxocariasis, birdshot choroidopathy, histoplasmosis (presumed ocular histoplasmosis syndrome), Behcet's

syndrome, sympathetic ophthalmia, Vogt-Koyanagi-Harada syndrome, sarcoidosis, reticulum cell sarcoma, large cell lymphoma, syphilis, tuberculosis, juvenile rheumatoid arthritis, endophthalmitis, and malignant melanoma of the choroid;

5

disorders of the retina, such as vascular retinopathies (e.g., arteriosclerotic retinopathy and hypertensive retinopathy), central and branch retinal artery occlusion, central and branch retinal vein occlusion, diabetic retinopathy (e.g., proliferative retinopathy and non-proliferative retinopathy), macular degeneration of the aged (age-related macular degeneration or senile macular degeneration), neovascular macular degeneration, retinal detachment, retinitis pigmentosa, retinal photic injury, retinal ischemia-induced eye injury, and glaucoma (e.g., primary glaucoma, chronic open-angle glaucoma, acute or chronic angle-closure, congenital (infantile) glaucoma, secondary glaucoma, and absolute glaucoma);

15

disorders of the optic nerve or visual pathways, such as papilledema (choked disk), papillitis (optic neuritis), retrobulbar neuritis, ischemic optic neuropathy, toxic amblyopia, optic atrophy, higher visual pathway lesions, disorders of ocular motility (e.g., third cranial nerve palsies, fourth cranial nerve palsies, sixth cranial nerve palsies, internuclear ophthalmoplegia, and gaze palsies);

25

free radical induced eye disorders and diseases;
and

immunologically-mediated eye disorders and
diseases, such as Graves' ophthalmopathy, conical
5 cornea, dystrophia epithelialis corneae, corneal
leukoma, ocular pemphigus, Mooren's ulcer, scleritis,
and sarcoidosis (See *The Merck Manual*, Sixteenth
Edition, 217:2365-2397 (1992) and *The Eye Book*,
Cassel, Billig, and Randall, The Johns Hopkins
10 University Press (1998)).

The compositions and methods of the present
invention are also useful in the treatment of the
following non-limiting eye injuries, and symptoms and
complications resulting therefrom: conjunctival and
15 corneal foreign body injuries, corneal abrasion,
intraocular foreign body injuries, lacerations, lid
lacerations, contusions, lid contusions (black eye),
trauma to the globe, laceration of the iris, cataract,
dislocated lens, glaucoma, vitreous hemorrhage,
20 orbital-floor fractures, retinal hemorrhage or
detachment, and rupture of the eyeball, anterior
chamber hemorrhage (traumatic hyphema), burns, eyelid
burns, chemical burns, chemical burns of the cornea
and conjunctiva, and ultraviolet light burns
25 (sunburn). See *The Merck Manual*, Sixteenth Edition,
217:2364-2365 (1992).

The compositions and methods of the present

invention are also useful in treating and/or preventing the following non-limiting exemplary symptoms and complications of eye disease, eye disorder or eye injury: subconjunctival hemorrhages, vitreous hemorrhages, retinal hemorrhages, floaters, retinal detachments, photophobia, ocular pain, scotomas (negative and positive), errors of refraction, emmetropia, ametropia, hyperopia (farsightedness), myopia (nearsightedness), astigmatism, anisometropia, aniseikonia, presbyopia, bleeding, recurrent bleeding, sympathetic ophthalmia, inflammation, swelling, redness of the eye, irritation of the eye, corneal ulceration and scarring, iridocyclitis, perforation of the globe, lid deformities, exophthalmos, impaired mobility of the eye, lid swelling, chemosis, loss of vision, including partial or total blindness, optic neuritis, fever, malaise, thrombophlebitis, cavernous sinus thrombosis, panophthalmitis, infection of the meninges and brain, papilledema, severe cerebral symptoms (headache, decreased level of consciousness, and convulsions), cranial nerve palsies, epiphora (chronic or persistent tearing), copious reflux of mucus or pus, follicular subconjunctival hyperplasia, corneal vascularization, cicatrization of the conjunctiva, cornea, and lids, pannus, hypopyon, lagophthalmos, phlyctenules, rubeosis iridis, bitemporal hemianopia, and homonymous

hemianopia. See *The Merck Manual, Sixteenth Edition*, 217:2362-2363 (1992).

The derivative may be administered in combination with an effective amount of one or more factor(s) useful in treating vision disorder, improving vision, treating memory impairment, or enhancing memory performance.

In a preferred embodiment, the factor(s) to be combined with the derivative is/are selected from the group consisting of immunosuppressants for treating autoimmune, inflammatory, and immunologically-mediated disorders; wound healing agents for treating wounds resulting from injury or surgery; antiglaucomatous medications for treating abnormally elevated intraocular pressure; neurotrophic factors and growth factors for treating neurodegenerative disorders or stimulating neurite outgrowth; compounds effective in limiting or preventing hemorrhage or neovascularization for treating macular degeneration; and antioxidants for treating oxidative damage to eye tissues.

Pharmaceutical Compositions of the Present Invention

25 The present invention also relates to a pharmaceutical composition comprising:

- (i) an effective amount of a derivative for treating a vision disorder, improving vision, treating memory impairment, or enhancing memory performance in an animal; and

- (ii) a pharmaceutically acceptable carrier.

5 The derivative may be administered in combination with an effective amount of one or more factor(s) useful in treating vision disorders, improving vision, treating memory impairment, or enhancing memory performance.





Table A
when D is a bond and R₂ is COOH,


	<u>No.</u>	<u>X</u>	<u>n</u>	<u>R₁</u>
5	1	O	1	3,4,5-trimethylphenyl
	2	O	2	3,4,5-trimethylphenyl
	3	O	1	tert-butyl
	4	O	3	tert-butyl
	5	O	1	cyclopentyl
10	6	O	2	cyclopentyl
	7	O	3	cyclopentyl
	8	O	1	cyclohexyl
	9	O	2	cyclohexyl
	10	O	3	cyclohexyl
15	11	O	1	cycloheptyl
	12	O	2	cycloheptyl
	13	O	3	cycloheptyl
	14	O	1	2-thienyl
	15	O	2	2-thienyl
20	16	O	3	2-thienyl
	17	O	1	2-furyl
	18	O	2	2-furyl
	19	O	3	2-furyl
	20	O	3	phenyl
25	21	O	1	1,1-dimethylpentyl
	22	O	2	1,1-dimethylhexyl
	23	O	3	ethyl

Table II.



No.	X	n	R_1	D	R_2
24	S	1	1,1-dimethyl propyl	CH_2	COOH
25	S	1	1,1-dimethyl propyl	bond	COOH
26	O	1	1,1-dimethyl propyl	CH_2	OH
27	O	1	1,1-dimethyl propyl	bond	SO_3H
28	O	1	1,1-dimethyl propyl	CH_2	CN
29	O	1	1,1-dimethyl propyl	bond	CN
30	O	1	1,1-dimethyl propyl	bond	tetrazolyl
31	S	1	phenyl	$(\text{CH}_2)_2$	COOH
32	S	1	phenyl	$(\text{CH}_2)_3$	COOH
33	S	2	phenyl	CH_2	COOH
34	O	1	1,1-dimethyl propyl	bond	CONH_2
35	O	2	1,1-dimethyl propyl	bond	CONH_2
36	S	2	2-furyl	bond	PO_3H_2
37	O	2	propyl	$(\text{CH}_2)_2$	COOH
38	O	1	propyl	$(\text{CH}_2)_3$	COOH
39	O	1	tert-butyl	$(\text{CH}_2)_4$	COOH
40	O	1	methyl	$(\text{CH}_2)_5$	COOH
41	O	2	phenyl	$(\text{CH}_2)_6$	COOH
42	O	2	3,4,5- trimethoxy-phenyl	CH_2	COOH
43	O	2	3,4,5- trimethoxy-phenyl	CH_2	tetrazolyl









TABLE III














	No.	n	X	D	R ₂	R ₁
5	44	1	S	bond	COOH	Phenyl
	45	1	O	bond	COOH	α-MethylBenzyl
	46	2	O	bond	COOH	4-MethylBenzyl
	47	1	O	bond	Tetrazole	Benzyl
	48	1	O	bond	SO ₃ H	α-MethylBenzyl
10	49	1	O	CH ₂	COOH	4-MethylBenzyl
	50	1	O	bond	SO ₂ HNMe	Benzyl
	51	1	O	bond	CN	α-MethylBenzyl
	52	1	O	bond	~PO ₃ H ₂	4-MethylBenzyl
	53	2	O	bond	COOH	Benzyl
15	54	2	O	bond	COOH	α-MethylBenzyl
	55	2	O	bond	COOH	4-MethylBenzyl
	56	2	S	bond	COOH	3,4,5-trimethoxy phenyl
	57	2	O	bond	COOH	Cyclohexyl
20	58	2	O	bond	PO ₂ HEt	i-propyl
	59	2	O	bond	PO ₃ HPropyl	ethyl
	60	2	O	bond	PO ₃ (Et) ₂	Methyl
	61	2	O	bond	OMe	tert-butyl
	62	1	O	bond	OEt	n-pentyl
25	63	2	O	bond	OPropyl	n-hexyl
	64	1	O	bond	OButyl	Cyclohexyl
	65	1	O	bond	OPentyl	cyclopentyl
	66	1	O	bond	OHexyl	n-heptyl
	67	1	O	bond	SMe	n-octyl
30	68	1	O	bond	SEt	n-nonyl
	69	2	O	bond	SPropyl	2-indolyl
	70	2	O	bond	SButyl	2-furyl
	71	2	O	bond	NHCOMe	2-thiazolyl
	72	2	O	bond	NHCOEt	2-thienyl
35	73	1	O	CH ₂	N(Me) ₂	2-pyridyl
	74	1	O	(CH ₂) ₂	N(Me)Et	1,1- dimethylpropyl
	75	1	O	(CH ₂) ₃	CON(Me) ₂	1,1- dimethylpropyl
40	76	1	O	(CH ₂) ₄	CONHMe	1,1- dimethylpropyl

	No.	n	X	D	R ₂	R ₁
5	77	1	O	(CH ₂) ₅	CONHEt	1,1-dimethylpropyl
	78	1	O	(CH ₂) ₆	CONHPropyl	1,1-dimethylpropyl
	79	1	O	bond	CONH(O)Me	Benzyl
10	80	1	O	bond	CONH(O)Et	α-Methylphenyl
	81	1	O	bond	CONH(O)Propyl	4-Methylphenyl
	82	1	O	(CH ₂) ₂	COOH	Benzyl
	83	1	O	bond	COOH	α-Methylphenyl
	84	1	O	bond	COOH	4-Methylphenyl
	85	1	O	CH ₂	COOH	1,1-dimethylpropyl
15	86	1	O	(CH ₂) ₂	COOH	1,1-dimethylbutyl
	87	1	O	(CH ₂) ₃	COOH	1,1-dimethylpentyl
20	88	1	O	(CH ₂) ₄	COOH	1,1-dimethylhexyl
	89	1	O	(CH ₂) ₅	COOH	1,1-dimethylethyl
	90	1	O	(CH ₂) ₆	COOH	iso-propyl
25	91	1	O	(CH ₂) ₇	COOH	tert-butyl
	92	1	O	(CH ₂) ₈	COOH	1,1-dimethylpropyl
30	93	1	O	(CH ₂) ₉	COOH	benzyl
	94	1	O	(CH ₂) ₁₀	COOH	1,1-dimethylpropyl
	95	1	O	C ₂ H ₂	COOH	cyclohexylmethyl
35	96	1	O	2-OH, Et	COOH	1,1-dimethylpropyl
	97	1	O	2-butylene	COOH	1,1-dimethylpropyl
40	98	1	S	i-Pro	COOH	1,1-dimethylpropyl
	99	2	S	tert-Bu	COOH	phenyl
	100	2	O	2-nitro-hexyl	COOH	1,1-dimethylpropyl
45	101	1	O	(CH ₂) ₂	CN	1,1-dimethylpropyl
	102	1	O	(CH ₂) ₃	CN	1,1-dimethylpropyl
50	103	3	O	bond	CONHNHSO ₂ Me	Benzyl
	104	3	O	bond	CONHNHSO ₂ Et	α-Methylphenyl
	105	3	O	bond	CONHSO ₂ Me	4-Methylphenyl
	106	1	O	bond	CONHNHSO ₂ Et	Phenyl

	No.	n	X	D	R2	R1
5	107	2	O	bond	CON (Me) CN	α -Methylphenyl
	108	1	O	bond	CON (Et) CN	4-Methylphenyl
	109	1	O	(CH ₂) ₂	COOH	methyl
	110	1	O	(CH ₂) ₃	COOH	ethyl
	111	1	O	(CH ₂) ₄	COOH	n-propyl
10	112	1	O	(CH ₂) ₅	COOH	t-butyl
	113	1	O	(CH ₂) ₆	COOH	Pentyl
	114	1	O	(CH ₂) ₇	COOH	Hexyl
	115	1	O	(CH ₂) ₈	COOH	Septyl
	116	1	O	(CH ₂) ₉	COOH	Octyl
	117	1	O	C ₂ H ₂	COOH	Cyclohexyl

No.	n	X	D	R2	R1
118	2	O	bond		1,1-dimethylpropyl
119	1	O	bond		1,1-dimethylpropyl
120	1	O	bond		1,1-dimethylpropyl
121	1	O	bond		1,1-dimethylpropyl
122	1	O	bond		1,1-dimethylpropyl
123	1	O	bond		1,1-dimethylpropyl
124	1	O	bond		1,1-dimethylpropyl
125	1	O	bond		1,1-dimethylpropyl

	No.	n	X	D	R2	R1
5	126	1	O	bond		1,1-dimethylpropyl
10	127	1	O	bond		1,1-dimethylpropyl
15	128	1	O	bond		1,1-dimethylpropyl
20	129	1	O	bond		1,1-dimethylpropyl
25	130	1	O	bond		1,1-dimethylpropyl
35	131	1	O	bond		1,1-dimethylpropyl
40	132	1	O	bond		1,1-dimethylpropyl
45	133	1	O	bond		1,1-dimethylpropyl

	No.	n	X	D	R2	R1
5	134	1	O	bond		1,1-dimethylpropyl
10	135	1	O	bond		1,1-dimethylpropyl
15						
20	136	1	O	bond		1,1-dimethylpropyl
25						
	137	1	O	bond	COOH	1,1-dimethylpropyl
30	138	2	O	bond	COOH	1,1-dimethylpropyl

Affinity for FKBP12

10 The compounds used in the inventive methods and
pharmaceutical compositions have an affinity for the
FK506 binding protein, particularly FKBP12. The
inhibition of the prolyl peptidyl *cis-trans* isomerase
activity of FKBP may be measured as an indicator of
15 this affinity.

K_i Test Procedure

Inhibition of the peptidyl-prolyl isomerase
(rotamase) activity of the compounds used in the
20 inventive methods and pharmaceutical compositions can
be evaluated by known methods described in the
literature (Harding et al., *Nature*, 1989, 341:758-760;
Holt et al. *J. Am. Chem. Soc.*, 115:9923-9938). These

values are obtained as apparent K_i 's.

The *cis-trans* isomerization of an alanine-proline bond in a model substrate, N-succinyl-Ala-Ala-Pro-Phe-
5 *p*-nitroanilide, is monitored spectrophotometrically in
a chymotrypsin-coupled assay, which releases *para*-
nitroanilide from the *trans* form of the substrate.
The inhibition of this reaction caused by the addition
of different concentrations of inhibitor is
10 determined, and the data is analyzed as a change in
first-order rate constant as a function of inhibitor
concentration to yield the apparent K_i values.

In a plastic cuvette are added 950 μ l of ice cold
assay buffer (25 mM HEPES, pH 7.8, 100 mM NaCl), 10 μ l
15 of FKBP (2.5 mM in 10 mM Tris-Cl pH 7.5, 100 mM NaCl,
1 mM dithiothreitol), 25 μ l of chymotrypsin (50 mg/ml
in 1 mM HCl) and 10 μ l of test compound at various
concentrations in dimethyl sulfoxide. The reaction is
initiated by the addition of 5 μ l of substrate
20 (succinyl-Ala-Phe-Pro-Phe-*para*-nitroanilide, 5 mg/ml
in 2.35 mM LiCl in trifluoroethanol).

The absorbance at 390 nm versus time is monitored
for 90 seconds using a spectrophotometer and the rate
constants are determined from the absorbance versus
25 time data files.

Route of Administration

To effectively treat vision loss or promote
45 vision regeneration, the compounds used in the
inventive methods and pharmaceutical compositions must

readily affect the targeted areas. For these purposes, the compounds are preferably administered [topically to the skin.]

[For topical application to the skin, the compounds can be formulated into suitable ointments containing the compounds suspended or dissolved in, for example, mixtures with one or more of the following: mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the compounds can be formulated into suitable lotions or creams containing the active compound suspended or dissolved in, for example, a mixture of one or more of the following: mineral oil, sorbitan monostearate, polysorbate 60, cetyl ester wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.]

Other routes of administration known in the pharmaceutical art are also contemplated by this invention.

Dosage

Dosage levels on the order of about 0.1 mg to about 10,000 mg of the active ingredient compound are useful in the treatment of the above conditions, with preferred levels of about 0.1 mg to about 1,000 mg. The specific dose level for any particular patient

will vary depending upon a variety of factors, including the activity of the specific compound employed; the age, body weight, general health, sex and diet of the patient; the time of administration; 5 the rate of excretion; drug combination; the severity of the particular disease being treated; and the form of administration. Typically, *in vitro* dosage-effect results provide useful guidance on the proper doses for patient administration. Studies in animal models 10 are also helpful. The considerations for determining the proper dose levels are well known in the art.

The compounds can be administered with other agents for treating vision loss, preventing vision degeneration, or promoting vision regeneration. 15 Specific dose levels for such other agents will depend upon the factors previously stated and the effectiveness of the drug combination.

5

10

15

The following examples are illustrative of preferred embodiments of the invention and are not to be construed as limiting the invention thereto. All polymer molecular weights are mean average molecular weights. All percentages are based on the percent by weight of the final delivery system or formulation prepared unless otherwise indicated and all totals equal 100% by weight.

20


EXAMPLES

25

The inventive compounds may be prepared by a variety of synthetic sequences that utilize established chemical transformations. A general pathway to the present compounds is described in Scheme I. N-glyoxylproline derivatives may be prepared by reacting L-proline methyl ester with methyl oxalyl chloride as shown in Scheme I.

The resulting oxamates may be reacted with a variety of carbon nucleophiles to obtain compounds used in the present invention.

Scheme I



EXAMPLE 1 (Compound 137)

Synthesis of (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate.

a. Synthesis of (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate.

A solution of L-proline methyl ester hydrochloride (3.08 g; 18.60 mmol) in dry methylene chloride was cooled to 0°C and treated with triethylamine (3.92 g; 38.74 mmol; 2.1 eq). After stirring the formed slurry under a nitrogen atmosphere for 15 min, a solution of methyl oxalyl chloride (3.20 g; 26.12 mmol) in methylene chloride (45 mL) was added dropwise. The resulting mixture was stirred at 0°C for 1.5 hr. After filtering to remove solids, the organic phase was washed with water,

dried over MgSO_4 and concentrated. The crude residue was purified on a silica gel column, eluting with 50% ethyl acetate in hexane, to obtain 3.52 g (88%) of the product as a reddish oil. Mixture of cis-trans amide rotamers; data for trans rotamer given. ^1H NMR (CDCl_3): δ 1.93 (dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H); 3.79, 3.84 (s, 3H total); 4.86 (dd, 1H, $J = 8.4, 3.3$).


b. Synthesis of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate.

A solution of methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90 mmol) in 30 mL of tetrahydrofuran (THF) was cooled to -78°C and treated with 14.2 mL of a 1.0 M solution of 1,1-dimethylpropylmagnesium chloride in THF. After stirring the resulting homogeneous mixture at -78°C for three hours, the mixture was poured into saturated ammonium chloride (100 mL) and extracted into ethyl acetate. The organic phase was washed with water, dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 2.10 g (75%) of the oxamate as a colorless oil. ^1H NMR (CDCl_3): δ 0.88 (t, 3H); 1.22, 1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H); 2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm, 1H, $J = 8.4, 3.4$).

c. Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-

2-pyrrolidinecarboxylic acid (Compound 137).

A mixture of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate (2.10 g; 8.23 mmol), 1 N LiOH (15 mL), and methanol (50 mL) was stirred at 0°C for 30 min and at room temperature overnight. The mixture was acidified to pH 1 with 1 N HCl, diluted with water, and extracted into 100 mL of methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification. ¹H NMR (CDCl₃): δ 0.87 (t, 3H); 1.22, 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H, J = 10.4, 7.3); 4.55 (dd, 1H, J = 8.6, 4.1).

Scheme IIEXAMPLE 2

Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxamide. (Compound 34)

Isobutyl chloroformate (20 mmol, 2.7 mL) was added to a solution containing (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid (4.89 g, 20 mmol) (from Example 1) in 50 mL methylene chloride at -10°C with stirring. After 5 minutes, ammonia was added dropwise (20 mmol, 10 mL of 2 M ethyl alcohol solution). The reaction was warmed up to room temperature after stirring at -10°C for 30 minutes. The mixture was diluted with water, and extracted into 200 mL methylene chloride. The organic extract was concentrated and further purified by silica gel to give 4.0 g of product

as a white solid (81.8% yield). ^1H NMR (CDCl_3): δ 0.91 (t, 3H, $J = 7.5$); 1.28 (s, 6H, each); 1.63-1.84 (m, 2H); 1.95-2.22 (m, 3H); 2.46 (m, 1H); 3.55-3.67 (m, 2H); 4.67 (t, 1H, $J = 7.8$); 5.51-5.53 (br, 1H, NH); 6.80 (br, 1H, NH).

EXAMPLE 3

Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile. (Compound 29)

To a solution of 0.465 mL DMF (6 mmol) in 10 mL acetonitrile at 0°C was added 0.48 mL (5.5 mmol) of oxalyl chloride. A white precipitate formed immediately and was accompanied by gas evolution. When complete, a solution of 1.2 g (5 mmol) of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxamide (from Example 2) in 2.5 mL acetonitrile was added. When the mixture became homogeneous, 0.9 mL (11 mmol) pyridine was added. After 5 min., the mixture was diluted into water and extracted by 200 mL ethyl acetate. The organic layer was concentrated and further purified by silica gel to give 0.3 g product as a white solid (72% yield). ^1H NMR (CDCl_3): δ 0.87 (t, 3H, $J = 7.5$); 1.22 (s, 3H); 1.24 (s, 3H); 1.80 (m, 2H); 2.03-2.23 (m, 4H); 3.55 (m, 2H); 4.73 (m, 1H).

EXAMPLE 4Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-
2-pyrrolidinetetrazole. (Compound 30)

A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-
2-pyrrolidinecarbonitrile (222 mg, 1 mmol) (from Example
3), NaN_3 (81 mg, 1.3 mmol) and NH_4Cl (70 mg, 1.3 mmol) in
3 mL DMF was stirred at 130°C for 16 hours. The mixture
was concentrated and purified by silica gel to afford 200
mg product as white solid (75.5% yield). ^1H NMR (CDCl_3):
 δ 0.88 (t, 3H, $J = 7.5$); 1.22 (s, 6H); 1.68 (m, 2H); 2.05-
2.36 (m, 3H); 2.85 (m, 1H); 3.54 (m, 1H); 3.75 (m, 1H);
5.40 (m, 1H).

5

10

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such modification are intended to be included within the scope of the following claims.

Example 5Synthesis of 3-phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (1)Methyl (2S)-1-(1,2-dioxo-2-methoxyethyl)-2-pyrrolidinecarboxylate

A solution of L-proline methyl ester
5 hydrochloride (3.08 g; 18.60 mmol) in dry methylene
chloride was cooled to 0°C and treated with
triethylamine (3.92 g; 38.74 mmol; 2.1 eq). After
stirring the formed slurry under a nitrogen atmosphere
for 15 min, a solution of methyl oxalyl chloride (3.20
10 g; 26.12 mmol) in methylene chloride (45 ml) was added
dropwise. The resulting mixture was stirred at 0°C for
1.5 hour. After filtering to remove solids, the
organic phase was washed with water, dried over MgSO₄
and concentrated. The crude residue was purified on
15 a silica gel column, eluting with 50% ethyl acetate in
hexane, to obtain 3.52 g (88%) of the product as a
reddish oil. Mixture of cis-trans amide rotamers;
data for trans rotamer given. ¹H NMR (CDCl₃): δ 1.93
(dm, 2H); 2.17 (m, 2H); 3.62 (m, 2H); 3.71 (s, 3H);
20 3.79, 3.84 (s, 3H total); 4.86 (dd, 1H, J = 8.4, 3.3).

Methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate

A solution of methyl (2S)-1-(1,2-dioxo-2-
methoxyethyl)-2-pyrrolidinecarboxylate (2.35 g; 10.90
25 mmol) in 30 ml of tetrahydrofuran (THF) was cooled to
-78°C and treated with 14.2 ml of a 1.0 M solution of
1,1-dimethylpropylmagnesium chloride in THF. After

stirring the resulting homogeneous mixture at -78°C for three hours, the mixture was poured into saturated ammonium chloride (100 ml) and extracted into ethyl acetate. The organic phase was washed with water, dried, and concentrated, and the crude material obtained upon removal of the solvent was purified on a silica gel column, eluting with 25% ethyl acetate in hexane, to obtain 2.10 g (75%) of the oxamate as a colorless oil. ^1H NMR (CDCl_3): d 0.88 (t, 3H); 1.22, 1.26 (s, 3H each); 1.75 (dm, 2H); 1.87-2.10 (m, 3H); 2.23 (m, 1H); 3.54 (m, 2H); 3.76 (s, 3H); 4.52 (dm, 1H, $J = 8.4, 3.4$).

Synthesis of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylic acid

A mixture of methyl (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarboxylate (2.10 g; 8.23 mmol), 1 N LiOH (15 ml), and methanol (50 ml) was stirred at 0°C for 30 minutes and at room temperature overnight. The mixture was acidified to pH 1 with 1 N HCl, diluted with water, and extracted into 100 ml of methylene chloride. The organic extract was washed with brine and concentrated to deliver 1.73 g (87%) of snow-white solid which did not require further purification. ^1H NMR (CDCl_3): d 0.87 (t, 3H); 1.22, 1.25 (s, 3H each); 1.77 (dm, 2H); 2.02 (m, 2H); 2.17 (m, 1H); 2.25 (m, 1H); 3.53 (dd, 2H, $J = 10.4, 7.3$); 4.55 (dd, 1H, $J = 8.6, 4.1$).

3-Phenyl-1-propyl (2S)-1-(3,3-dimethyl-1,2-dioxopentyl)-2-pyrrolidinecarboxylate (1)

A mixture of (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidine-carboxylic acid (600 mg; 2.49 mmol), 3-phenyl-1-propanol (508 mg; 3.73 mmol), dicyclohexylcarbodiimide (822 mg; 3.98 mmol), camphorsulfonic acid (190 mg; 0.8 mmol) and 4-dimethylaminopyridine (100 mg; 0.8 mmol) in methylene chloride (20 ml) was stirred overnight under a nitrogen atmosphere. The reaction mixture was filtered through Celite to remove solids and concentrated in vacuo, and the crude material was purified on a flash column (25% ethyl acetate in hexane) to obtain 720 mg (80%) of Example 1 as a colorless oil. ¹H NMR (CDCl₃): δ 0.84 (t, 3H); 1.19 (s, 3H); 1.23 (s, 3H); 1.70 (dm, 2H); 1.98 (m, 5H); 2.22 (m, 1H); 2.64 (m, 2H); 3.47 (m, 2H); 4.14 (m, 2H); 4.51 (d, 1H); 7.16 (m, 3H); 7.26 (m, 2H).

Figure 1. GPI 1046 protects retinal ganglion cells against degeneration following retinal ischemia.

Retinal ganglion cells were retrogradely labeled in adult rats by bilateral injection of fluorogold in their lateral geniculate nuclei. Labeled ganglion cells in the normal rat retina appear as white profiles against the dark background (Figure 1A). Complete retinal ischemia was produced by infusing normal saline solution into the retinal vitreous cavity of each eye until the intraocular pressure exceeded arterial blood pressure. 28 days after the ischemic episode extensive degeneration of retinal ganglion cell was evidenced by massive reduction in the density of fluorogold labeled cells (Figure 1B). Administration of GPI 1046 (10mg/kg, s.c.) 1 hour prior to the ischemic episode and at 10mg/kg/day for the next four days produced noticeable protection of a large proportion of the vulnerable ganglion cell population (Figure 1C).

Figure 2. GPI 1046 prevents degeneration of optic nerve axons and myelin following retinal ischemia

Examination of the optic nerves from the same retinal ischemia cases reveals that GPI 1046 produces dramatic protection of optic nerve element from ischemic degeneration. Toluidine blue staining of epon embedded optic nerve cross sections revealed the detail of myelin sheaths (white circles) and optic nerve axons

(black centers) in the normal rat optic nerve. Optic nerves from vehicle treated cases examined 28 days after a 1 hour retinal ischemic episode are characterized by a decreased density of optic nerve axons and the appearance of numerous degenerating myelin figures (bright white filled circles). Treatment with GPI 1046 protected the majority of optic nerve axons from degeneration and also dramatically decreased the density of degenerating myelin figures.

Figure 3. GPI 1046 provides moderate protection against retinal ganglion cell death after optic nerve transection

Complete transection of the optic nerve 5 mm from the eyeball produces massive degeneration of retinal ganglion cells, representing loss of >87% of the normal ganglion cell population 90 days after the injury (Table 1). Few spared fluorogold pre labeled ganglion cells are present in vehicle treated cases (large white figures) among a population of small microglia that digest the debris of the degenerating cells and take up the fluorogold label (Figure 3A). Treatment with GPI 1046 for 14 days resulted in a small but not significant increase in the density of retinal ganglion cells that survived 90 days after transection (Table 1) but treatment with GPI 1046 for the first 28 days after transection produced moderate but significant

protection of 12.6% of the vulnerable ganglion cell population (Table 1, Figure 3B).

Figure 4. GPI 1046 treatment duration significantly affects the process of optic nerve axonal degeneration after transection.

Examination of optic nerve axon density in the proximal stump of the optic nerve from the same cases revealed a more dramatic protection afforded by GPI 1046 treatment. 90 days after transection few ganglion cell axons remain within the optic nerve (Figure 4B), representing only 5.6% of the normal population. The loss of axons reflects both the death of retinal ganglion cells and the regression or "dying back" of the axons of ~ 70% of the small surviving ganglion cell population into the retina itself (Table 1). Treatment with GPI 1046 for the first 14 days after optic nerve transection produced a small but significant 5.3% protection of optic nerve axons (Figure 4D, Table 1), but treatment with the same dose of GPI 1046 for 28 days resulted in the protection of optic nerve axons for the vast majority (81.4%) of spared retinal ganglion cells (Figure 4C, Table 1).

Figure 5. GPI 1046 treatment produces a greater effect on optic nerve axons than ganglion cell bodies

This summary figure shows data from Figure 3 ganglion cell protection and higher power photomicrographs of

optic nerve axon protection (Figure 5A&B, upper panels). 28 day treatment with GPI 1046 produced a significant increase in the density of large, and particularly medium and small caliber optic nerve axons (Figure 5C&D, lower panels).

Figure 6. GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in the proximal stump

Myelin basic protein immunohistochemistry labels fascicles (darker labeled 'islands') of myelinated axons in the normal optic nerve (Figure 6A, upper left). 90 days after transection extensive degeneration of myelin is evident in vehicle treated cases, characterized by the loss of fascicular organization and the appearance of numerous large dense degenerating myelin figures (Figure 6B, upper right). Treatment with GPI 1046 for the first 14 days after optic nerve transection did not alter the pattern of myelin degeneration (Figure 6C, lower left panel), and yielded an insignificant 1.6% quantitative recovery in myelin density (Table 1). Extending the GPI 1046 treatment course through the first 28 days after optic nerve transection produced a dramatic preservation of the fascicular staining pattern for myelin basic protein in the proximal stump of the optic nerve and decreased the density of degenerating myelin figures

(Figure 6D, lower right panel), representing a '70% recovery of myelin density (Table 1).

Figure 7. FKBP-12 immunohistochemistry labels

5 oligodendroglia (large dark cells with fibrous processes), the cells which produce myelin, located between the fascicles of optic nerve fibers, and also some optic nerve axons.

10 **Figure 8. GPI 1046 treatment for 28 days after optic nerve transection prevents myelin degeneration in the distal stump.**

Complete transection of the optic nerve leads to degeneration of the distal segments (axon fragments disconnected from the ganglion cell bodies), and the
15 degeneration of their myelin sheaths. 90 days after transection (Figure 8B) myelin basic protein immunohistochemistry reveals the near total loss of fascicular organization (present in the normal optic
20 nerve, Figure 8A) and the presence of numerous dense degenerating myelin figures. Quantitation reveals that the cross sectional area of the transected distal stump shrinks by 31% and loses approximately 1/2 of its myelin (Table 1). Treatment with GPI 1046 for the
25 first 14 days after transection did not protect against shrinkage of the distal stump but did slightly increase the density of myelin, though the density of

degenerating myelin figures remained high (Figure 8C, Table 1). GPI 1046 treatment through the first 28 days produced dramatic protection of the fascicular pattern of myelin labeling, decreased the density of

5 degenerating myelin figures, prevented cross sectional shrinkage of the distal stump of the transected nerve and maintained the myelin levels at ~99% of normal levels (Figure 8D, Table 1).

10 **Figure 9. 28 day treatment with GPI 1046 treatment beginning 8 weeks after onset of streptozotocin induced diabetes decreases the extent of neovascularization in the inner and outer retina and protects neurons in the inner nuclear layer (INL) and ganglion cell layer (GCL) from degeneration.**

15 Negative images of cresyl violet stained tangential retinal sections reveals perikarya in the three cellular layers (Figure 9A). The retinae of streptozotocin treated animals administered only
20 vehicle (Figure 9B) exhibited loss of cells from the ONL and INL, decreased thickness of the Outer plexiform layer (the dark area between ONL and INL) and a dramatic increase in the size and density of retinal blood vessels (large black circular outlines) in the
25 INL, OPL, ONL and the photoreceptor layer (PR, the gray fuzzy area above the ONL). GPI 1046 treatment reduced neovascularization (i.e. prevented the proliferation of

blood vessels) in the PR, ONL, OPL and INL. Although
GPI 1046 did not appear to protect against neuronal
loss in the ONL, it appeared to decrease the loss of
neurons in both the INL and GCL compared to
5 streptozotocin/vehicle treated controls.

Example 6

10

In Vivo Retinal Ganglion Celland Optic Nerve Axon Tests

15

The extent of degeneration reduction or prevention in retinal ganglion cells and optic nerve axons was determined in a vision loss model utilizing surgical optic nerve transection to simulate mechanical damage to the optic nerve. The effects of several neuroimmunophilin FKBP ligands on retinal ganglion cells neuroprotection and optic nerve axon density was determined experimentally, comparing 14 day and 28 day neuroimmunophilin FKBP ligand treatments. The effects of treatment with neuroimmunophilin FKBP ligands on retinal ganglion cells and optic nerve axons was correlated.

20

Surgical Procedures

25

Adult male Sprague Dawley rats (3 months old, 225-250 grams) were anesthetized with a ketamine

(87mg/kg) and xylazine (13mg/kg) mixture. Retinal ganglion cells were pre-labeled by bilateral stereotaxic injection of the fluorescent retrogradely transported marker fluoro-gold (FG, 0.5 microliters of 2.5% solution in saline) at the coordinates of the LGNd (4.5 millimeters post β , 3.5 millimeters lateral, 4.6 millimeters below dura). Four days later, FG labeled rats underwent a second surgery for microsurgical bilateral intraorbital optic nerve transection 4-5 millimeters behind the orbit.

Experimental animals were divided into six experimental groups of six rats (12 eyes) per group. One group received a neuroimmunophilin FKBP ligand (10 milligrams per kg per day sc in PEG vehicle (20 percent propylene glycol, 20 percent ethanol, and 60 percent saline)) for 14 days. A second group received the same neuroimmunophilin FKBP ligand dose for 28 days. Each treated group had a corresponding sham/surgery and transection control group which received corresponding 14 or 28 day dosing with the vehicle only.

All animals were sacrificed 90 days after optic nerve transection and perfused pericardially with formalin. All eyes and optic nerves stumps were removed. Cases were excluded from the study if the optic nerve vasculature was damaged or if FG labeling

was absent in the retina.

Retinal Ganglion Cell Counts

Retinas were removed from eyes and prepared for wholemount analysis. For each group, five eyes with
5 dense and intense FG labeling were selected for quantitative analysis using a 20 power objective. Digital images were obtained from five fields in the central retina (3-4 millimeters radial to optic nerve head). FG labeled Large ($>18\text{ }\mu\text{m}$), medium ($12\text{-}16\text{ }\mu\text{m}$),
10 and small ($<10\text{ }\mu\text{m}$) ganglion cells and microglia were counted in five $400\text{ }\mu\text{m}$ by $400\text{ }\mu\text{m}$ fields per case, 5 cases per group.

Examination of Optic Nerves

Proximal and distal optic nerve stumps were
15 identified, measured, and transferred to 30% sucrose saline. The proximal stumps of five nerves were blocked and affixed to a chuck, and 10 micron cross sections were cut on a cryostat; one in ten sections were saved per set. Sections including the region 1-2
20 mm behind the orbit were reacted for RT97 neurofilament immunohistochemistry. Analysis of optic nerve axon density was performed using a 63 power oil immersion lens, a Dage 81 camera, and the Simple Image Analysis program. RT97 positive optic nerve axons
25 were counted in three $200\text{ }\mu\text{m}$ by $200\text{ }\mu\text{m}$ fields per nerve. The area of the nerve was also determined for

each case at 10 power.

As depicted graphically in Table I&II, the 14 day course of treatment with a neuroimmunophilin FKBP ligand provided moderate neuroprotection of retinal ganglion cells observed 28 days after optic nerve transection. However, by 90 days after transection, only 5% of the ganglion cell population remained viable.

90 days after optic nerve transection the number of axons persisting in the proximal stump of the optic nerve represented approximately one half of the number of surviving ganglion cells in groups of animals that received vehicle alone or the 14 day course of treatment with a neuroimmunophilin FKBP ligand. These results indicate that over half of the transected ganglion cell axons retract beyond the optic nerve head, and that treatment with a neuroimmunophilin FKBP ligand during the first 14 days after optic nerve transection is not sufficient to arrest this retraction.

As depicted graphically in Table I&II,, more prolonged treatment with a neuroimmunophilin FKBP ligand during the 28 day course of treatment produced a moderate increase in retinal ganglion cell neuroprotection. Approximately 12% of the vulnerable retinal ganglion cell population was protected. A

Table 1.
Effect of prolonged GPI 1046 treatment on retinal ganglion cell survival,
optic nerve axon preservation, and myelination 90 days after optic nerve transection

GROUP	RGC Counts ¹	ON Axon density ²	ON head area (%sham)	% RGCs Rescued	increased ON axon density ³	Spared RGC population	ON axon Count ⁴	% surviving RGCs with ON axons	Proximal optic nerve myelin basic protein Density ⁵	Distal optic nerve myelin basic protein Density ⁵
Sham	290 ± 14.8	7600*	100%	-		120,000*	120,000	100%	normal	
ONT/Vehicle	35.9 ± 2.8	428 ± 34	68%	(87% loss)		14,855	4593	30.9%	52 ± 5.2 SEM % loss	31% shrinkage 52.1% loss
ONT/ 14 days GPI 1046	49 ± 5.3	569 ± 23	76%	5.3%	1.5X	20,275	6820	33.6%	1.6 ± 3.0 SEM % recovery	33% shrinkage 47% loss
ONT/ 28 days GPI 1046	67.9 ± 5.8*	1526 ± 120*	95%*	12.6%*	5.0X	28,096*	22,861*	81.4%	70 ± 6.3 SEM % recovery*	56% less shrinkage* 99% myelin preservation*

*significance p<.001

¹ Mean density + SEM of Fluoro-gold labeled retinal ganglion cells (RGC) in 400 µm x 400 µm sample gridfields.

² mean density + SEM of RT97 neurofilament antibody labeled optic nerve (ON) axons in 200 µm x 200µm region of interest

*estimate for 200 µm x 200µm region in normal optic nerve assuming 120,000 RGC axons in normal rat optic nerve, measured to be 0.630 mm² mean cross sectional area

³adjusted for optic nerve diameter


⁴ calculated by multiplying axonal density by ON area

⁵ determined from 20X analysis of % areal coverage of optic nerve cross section

⁶shrinkage determined by comparing cross sectional area to sham control, myelin levels determined by multiplying cross sectional area by myelin density

TABLE II

**Neuroprotective effect of GPI 1046
on retinal ganglion cells following optic nerve transection**



similar proportion (~50%) of optic nerve axon density sparing was also observed. These results demonstrate the startling result that extending the duration of treatment with a neuroimmunophilin FKBP ligands to 28
5 days after transection completely arrests the regression of damaged axons for essentially the entire surviving population of retinal ganglion cells.

Additional results are set forth in Tables III & IV.

TABLE III

Correlation between Retinal Ganglion Cell and Optic Nerve Axon Sparing at 90 days following optic nerve transection and 14 or 28 day GPI 1046 treatment





TABLE IV

GPI 1046 preserves optic nerve axons
in the proximal stump following transection



Row Number

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

Example 7

A patient is suffering from macular degeneration.

A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 8

A patient is suffering from glaucoma, resulting in cupping of the optic nerve disc and damage to nerve fibers. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 9

A patient is suffering from cataracts requiring surgery. Following surgery, a derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical

composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 10

A patient is suffering from an impairment or blockage of retinal blood supply relating to diabetic retinopathy, ischemic optic neuropathy, or retinal artery or vein blockage. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 11

A patient is suffering from a detached retina. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or

promotion of vision regeneration are/is expected to occur following treatment.

Example 12

5 A patient is suffering from tissue damage caused
by inflammation associated with uveitis or
conjunctivitis. A derivative as
identified above, alone or in combination with one or
more other neoplastic factors, or a pharmaceutical
10 composition comprising the same, may be administered
to the patient. A reduction in vision loss,
prevention of vision degeneration, and/or promotion of
vision regeneration are/is expected to occur following
treatment.

15

Example 13

 A patient is suffering from photoreceptor damage
caused by chronic or acute exposure to ultraviolet
light. A derivative as identified above,
20 alone or in combination with one or more other neoplastic
factors, or a pharmaceutical composition comprising
the same, may be administered to the patient. A
reduction in vision loss, prevention of vision
degeneration, and/or promotion of vision regeneration
25 are/is expected to occur following treatment.

Example 14

A patient is suffering from optic neuritis. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 15

A patient is suffering from tissue damage associated with a "dry eye" disorder. A derivative as identified above, alone or in combination with one or more other neoplastic factors, or a pharmaceutical composition comprising the same, may be administered to the patient. A reduction in vision loss, prevention of vision degeneration, and/or promotion of vision regeneration are/is expected to occur following treatment.

Example 16

**Efficacy of representative compounds from
different immunophilin ligand series
in protecting retinal ganglion cell axons from
degeneration following optic nerve transection
is set forth in Table V.**

**Efficacy of representative compounds from
different immunophilin ligand series
in protecting retinal ganglion cell axons from
degeneration following optic nerve transection**







Compound	Structure	Comments	RT 27-29°C axon density 14d post-ON transection (% ON non-transected)
B		Adamanti Thioester of urea Ki rotomase = 149 nM Clearance = ? μ l/min	100.0% \pm 5.2% SEM
A GPI 1046		Ester Ki rotomase = 7.5 nM Clearance = 63.8 μ l/min	60.5% \pm 3.9 SEM
C		Sulfonamide Ki rotomase = 107 nM Clearance = 31.1 μ l/min	60.4% \pm 3.1 % SEM
D		Pipecolic sulfonamide Ki rotomase = nM Clearance = μ l/min	58.4% \pm 6.4% SEM
E		Ester of pipecolic acid Ki rotomase = 20 nM Clearance = 41.8 μ l/ml	56.6% \pm 9.4% SEM
F		Proline heterocycle Analog of GPI 1046 Ki rotomase = 272 nM Clearance = ? μ l/min	55.1% \pm 5.9% SEM

TABLE V









G		Pipercolic acid dimethyl ketone Ki rotomase >10,000 nM Clearance=? $\mu\text{l/min}$	34.0% $\pm 4.8\%$ SEM
H		Ki rotomase = nM Clearance=? $\mu\text{l/min}$	30.3% $\pm 8.0\%$ SEM
I		Ester of Thiourea Ki rotomase= 131 nM Clearance=8.0 $\mu\text{l/min}$	23.8% ± 5.3 SEM
J		Ketone analog of GPI 1046 Ki rotomase= 210nM Clearance=1.5 $\mu\text{l/min}$	15.8% $\pm 4.8\%$ SEM
K		Pipercolic acid Thioester Ki rotomase= 86nM Clearance= 4.5 $\mu\text{l/min}$	13.0% $\pm 4.2\%$ SEM
L		Prolyl acid Ki rotomase= >7743nM Clearance=5.2 $\mu\text{l/min}$	7.8% $\pm 3.0\%$ SEM
M		Thioester Ki rotomase =7nM Clearance=12.5 $\mu\text{l/min}$	-6.3% $+3.9\%$ SEM
N		Ki rotomase = 722 nM Clearance= 21.9 $\mu\text{l/ml}$	

TABLE V continued

Example 17

5 **THE FKBP NEUROIMMUNOPHILIN LIGAND GPI-1046**
 ENHANCES RETINAL GANGLION CELL SURVIVAL
 AND ARRESTS AXONAL DYING BACK
 FOLLOWING OPTIC NERVE TRANSECTION

10

Transection of the mammalian optic nerve results in a brief period of abortive regeneration, but the majority of axotomized neurons die and the axons from many persisting ganglion cells die back beyond the optic nerve head. The present Example was designed to examine the neuroprotective effects of GPI-1046 following optic nerve transection.

Retinal ganglion cells in adult male Sprague
20 Dawley rats were retrogradely labeled by fluorogold injection in the LGNd and four days later the optic nerves were transected 5 mm behind the globe. Groups of animals received either GPI-1046 10mg/kg/day s.c. or vehicle for 28 days. All experimental animals and
25 controls were sacrificed 90 days after transection.

By 90 days only - 10% of the FG labeled ganglion cell population survived but less than half of these neurons maintained axons that extended past the optic
30 nerve head, as detected with RT97 neurofilament immunohistochemistry. GPI-1046 treatment produced a moderate degree of perikaryal neuroprotection, sparing

25% of the ganglion cell population, and preserved the axons of virtually all protected neurons in the proximal stump of the transected nerve. These results indicate that treatment with the FKBP neuroimmunophilin ligand GPI-1046 produces a fundamental alteration in the pathological process following injury to CNS tracts.

These results also demonstrate that the small molecule FKBP neuroimmunophilin ligand GPI 1046 enhances neurite outgrowth in culture, enhance peripheral nerve regeneration, and stimulate sprouting within the CNS following partial deafferentation.

Example 18

5 NEUROIMMUNOPHILIN LIGANDS PROMOTE RECOVERY
FROM THE PERIPHERAL SENSORY NEUROPATHY ASSOCIATED
WITH STREPTOZOTOCIN-INDUCED DIABETES

10 Peripheral neuropathy is a common debilitating
complication of Type 2 diabetes in some 30-40% of
diabetic patients. Neurotrophic factors such as nerve
growth factor (NGF) are known to promote survival of
developing and adult neurons of the peripheral nervous
15 system (PNS), and have also been evaluated as
treatments for diabetic peripheral neuropathy. Some of
the selective ligands of the neuroimmunophilin FKBP-12
such as the small molecule GPI-1046, have also been
shown to promote repair and regeneration in the central
20 and peripheral nervous systems (Proc. Nat'l. Acad. Sci.
USA 94 , 2019-2024, 1997).

 In this Example the potential therapeutic effects
of GPI-1046 were evaluated for its ability to improve
25 sensory function in the streptozotocin-induced diabetic
rat. The procedure involved using Male Wistar rats
which were given a single injection of streptozotocin
(65 mg/kg i.v.). Blood glucose levels were determined
weekly for the first three weeks and on the last week
30 of the experiment. Animals were evaluated weekly for
signs of sensory neuropathy using the conventional hot
plate and tail flick apparatus test procedures. After

six weeks, treatment either with GPI-1046 or vehicle was initiated.

The results demonstrated that behavioral testing using the hot plate and the tail flick apparatus indicated improvement in latency in lesioned animals treated for 6 weeks with GPI-1046 at 10 mg/kg s.c. The results also showed that GPI-1046 ameliorates the behavioral sequelae of diabetic sensory neuropathy and may offer some relief for patients suffering from diabetic peripheral neuropathy.

Morris Watermaze/Aging and Memory Test Procedure

5

Aged rodents exhibit marked individual differences in performance on a variety of behavioral tasks, including two-choice spatial discrimination in a modified T-maze, spatial discrimination in a circular platform task, 10 passive avoidance, radial maze tasks, and spatial navigation in a water pool.

15

In all of these tasks, a proportion of aged rats or mice perform as well as the vast majority of young control animals, while other animals display severe impairments in memory function compared to young animals. For example, Fischer and colleagues showed that the proportion of rats displaying significant impairments in spatial navigation increases with age, (Fischer et al. 20 1991b) with 8% of all 12 month old, 45% of 18 month old, 53% of 24 month old, and 90% of all 30 month old rats displaying impairments in spatial acquisition of the Morris watermaze task relative to young controls.

25

Specifically, rodent spatial learning and memory decline during aging has been accepted by many investigators as an intriguing correlative animal model of human senile dementia. Cholinergic function in the hippocampus has

been extensively studied as a component of spatial learning in rodents, and declining hippocampal cholinergic function has been noted in parallel with the development of learning and memory impairments. In addition, other neurotransmitter systems have been shown to contribute to spatial learning, and to decline with age, such as the dopaminergic and noradrenergic, serotonergic, and glutamatergic systems.

Also, reports on age-related deficits of hippocampal long-term potentiation (LTP)-induction, a reduction in theta rhythm frequency, a loss of experience-dependent plasticity of hippocampal place-units, and reductions in hippocampal protein kinase C are in keeping with the concept that no single underlying pathology can be identified as the cause of age-related behavioral impairment in rodents. However, the various experimental therapeutic approaches that have been undertaken to improve memory function in aged rodents have been somewhat slanted towards the cholinergic hypothesis.

The Morris watermaze is widely used for assessing spatial memory formation and retention in experimental animals. The test depends on the animal's ability to utilize spatial visual information in order to locate a submerged escape platform in a water tank. It is important that the tank itself be as devoid of specific visual features

as possible - thus, it is always circular in shape, the sides are kept smooth and in uniform dull colors, and the water is rendered opaque with nontoxic watercolour pigment or powdered milk. This is to ensure that the animal navigates only by the use of more distant visual cues, or by the use of intra-maze cues specifically provided by the experimenter.

The tank is filled to a level which forces the animal to swim actively. Normal mice and rats react aversively to the swimming part of the test and will climb onto, and remain on, an escape platform from which they are removed to a heated resting cage.

If the platform is visible (i.e. above the surface), animals placed in the tank will quickly learn to home in on the platform and climb out onto it. Testing with a visible platform will also ensure that the experimental animals are not blind and show sufficient motivation and stamina to perform the task, which can be important in experiments involving aged rodents. If the platform is invisible (i.e. submerged just below the surface), normal animals learn to use distant visual cues in the test room for orientation in the test tank, and, when placed in the tank, will quickly home in on the approximate location of the platform and circle in that area until the platform is found.

The animals' path, speed, and swim time are tracked with a ceiling camera for later computerized analysis. Over the course of several successive trials, spatial learning can therefore be defined as a drop of distance swum, or time elapsed, from placement in the tank until escape onto the invisible platform.

The test can be adapted to assess several aspects of spatial memory: a) acquisition of a cued task, where the animal's ability to link one visual cue directly with the escape platform depends on cortical function (i.e. a ball is suspended over the escape platform and the animal learns to follow this cue to find the platform); b) acquisition of a spatial task, where the animal's ability to learn the location of a submerged escape platform based on a combination of distant visual cues is dependent upon hippocampal function (i.e. the animal learns to triangulate its position in the tank by visually aligning the paper-tower dispenser with the door and ceiling lamp); c) retention of a successfully acquired spatial task, which is predominantly dependant on cortical function (i.e. the animal must remember the spatial location of the platform over several weeks); d) a hippocampus-dependant reversal task where the animals must reacquire a new spatial platform location (i.e. the platform is moved to a new location between swim trials

and the animal must abandon its previous search strategy and acquire a new one).

5 These different modifications of the Morris watermaze procedure can be applied in sequence to the same set of experimental animals and allow for a thorough characterization of their spatial memory performance and its decline with normal ageing. Moreover, such a series of sequential memory tests sheds some light on the functional integrity of the specific brain systems involved in the acquisition and retention of spatial memory (e.g. rats with cholinergic lesions of the hippocampus may remember a platform location acquired weeks before, but persevere over the old platform location after the platform is moved).

10

15

Example 19

EFFECTS OF CHRONIC GPI-1046 ADMINISTRATION ON SPATIAL LEARNING AND MEMORY IN AGED RODENTS

20

This Example shows the effects of chronic treatment with the systemically available FKBP-ligand GPI-1046 on spatial learning and memory in aged rodents.

25

The procedure involved using three-month old (young) and 18-19 month old male C57BL/6N-Nia (aged) mice which

habituated to the well known and conventional Morris watermaze during a 4 trials/day, 3-4 day visible platform training phase. Subsequent spatial acquisition testing was conducting as follows: All mice were given 4
5 trials/day (block), for 5 days. Maximum swim time was 90 seconds. Aged mice were allocated to an "aged impaired" group if their performance during blocks 4 or 5 of the acquisition phase was >1 S.D. above the mean of "young" mice, and to an "aged non-impaired" group if their
10 performance was < 0.5 S.D. above the mean of "young" mice. Aged groups were then split into statistically similar "GPI-1046" and "vehicle" groups.

Daily treatment with 10mg/kg GPI-1046 was initiated
15 3 days after the end of acquisition training, and continued through retention testing. Retention testing began after 3 weeks of dosing using the same methods as the acquisition phase. Swim Distances (cm) were analyzed in a 7 X 5 ANOVA including Groups and Blocks (1-5) as
20 factors in the analysis, treating Blocks as a repeated measure.

The results showed that planned contrasts revealed that there were significant differences between the
25 "young", and "aged impaired-vehicle and GPI-1046" treated groups at the end of the acquisition phase, $F_{1,58} = 26.75$, $P=0.0001$, and $F_{1,58} = 17.70$, $P=0.0001$ respectively. While

there were no significant differences between the two "aged impaired" groups, $F_{1,58} = 0.67$, $P = 0.42$. During retention testing, however, "aged impaired-vehicle" treated animals performed significantly poorer than "aged impaired - GPI-1046", and "young" animals, $F_{1,69} = 8.11$, $P = 0.006$, and $F_{1,69} = 25.45$, $P = 0.0001$ respectively. There was no longer any statistically significant difference between the "young" and "aged impaired" - GPI-1046" treated groups during the retention phase, $F_{1,69} = 3.09$, $P = 0.08$. In summary, systemic treatment with GPI-1046 significantly enhanced spatial memory performance of mice with age-related spatial memory impairments.

The invention being thus described, it will be obvious that the same may be varied in many ways. Such variations are not to be regarded as a departure from the spirit and scope of the invention and all such
5 modification are intended to be included within the scope of the following claims.

What is claimed is:

1. A method for treating a vision disorder,
5 improving vision, treating memory impairment or enhancing
memory performance in an animal, which comprises
administering to said animal an effective amount of an N-
heterocyclic ring compound containing a carboxylic acid
or carboxylic acid isostere moiety thereof attached to
10 the 2-carbon of the N-heterocyclic ring.
2. The method of claim 1, wherein the N-
heterocyclic ring compound is immunosuppressive or non-
immunosuppressive.
- 15 3. The method of claim 1, wherein the N-
heterocyclic ring compound has an affinity for an FKBP-
type immunophilin.
- 20 4. The method of claim 3, wherein the FKBP-type
immunophilin is FKBP-12.
5. The method of claim 1, wherein the vision
disorder is selected from the group consisting of: visual
25 impairments; orbital disorders; disorders of the lacrimal
appartus; disorders of the eyelids; disorders of the

5

10

15



n is 1-3;

R₁ is selected from the group consisting of C₁-C₉,




















20

straight or branched chain alkyl, C₂-C₉ straight or branched chain alkenyl, aryl, heteroaryl, carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl; and
R₂ is a carboxylic acid or a carboxylic acid isostere; or a pharmaceutically acceptable salt, ester, or solvate thereof.

8. The method of claim 7, wherein R₂ is a carbocycle or heterocycle containing any combination of CH₂, O, S, or N in any chemically stable oxidation state, where any of the atoms of said ring structure are optionally substituted in one or more positions with R³, wherein
R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched chain alkyl, C₂-C₆ straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched chain alkyl or alkenyl.

9. The method of claim 7, wherein R₂ is selected from the group below:



where the atoms of said ring structure R_2 may be optionally substituted at one or more positions with R^3 , wherein

R^3 is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl, alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy, cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl, thioalkyl, alkylthio, sulfonyl, C_1 - C_6 straight or branched chain alkyl, C_2 - C_6 straight or branched chain alkenyl or alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and CO_2R^4 where R^4 is hydrogen or C_1 - C_9 straight or branched chain alkyl or alkenyl.

10. The method of claim 7, wherein R_2 is selected from the group consisting of $-COOH$, $-SO_3H$, $-SO_2HNR^3$, $-PO_2(R^3)_2$, $-CN$, $-PO_3(R^3)_2$, $-OR^3$, $-SR^3$, $-NHCOR^3$, $-N(R^3)_2$, $-CON(R^3)_2$, $-CONH(O)R^3$, $-CONHNHSO_2R^3$, $-COHNSO_2R^3$, and $-CONR^3CN$.

11. The method of claim 7, wherein the N-heterocyclic ring compound is selected from the group consisting of: (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-hydroxymethyl pyrrolidine; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinetetrazole; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-pyrrolidinecarbonitrile; (2S)-1-(1,2-dioxo-3,3-dimethylpentyl)-2-aminocarbonyl piperidine; and compounds 1-25, 27, 28, 31-33, and 35-136

of Tables I, II, and III.

12. A pharmaceutical composition for treating a vision disorder, improving vision, treating memory
5 impairment or enhancing memory performance in an animal, comprising:

- 10 a) an effective amount of an N-heterocyclic carboxylic acid or carboxylic acid isostere for treating a vision disorder, improving vision, treating memory impairment or enhancing memory performance in an animal; and
- b) a pharmaceutically acceptable carrier.

15 13. The pharmaceutical composition of claim 12, wherein the N-heterocyclic carboxylic acid or carboxylic acid isostere is immunosuppressive or non-immunosuppressive.


20 14. The pharmaceutical composition of claim 12, wherein the N-heterocyclic ring compound has an affinity for an FKBP-type immunophilin.

25 15. The pharmaceutical composition of claim 14, wherein the FKBP-type immunophilin is FKBP-12.

16. The pharmaceutical composition of claim 12,
wherein the vision disorder is selected from the group
consisting of: visual impairments; orbital disorders;
disorders of the lacrimal apparatus; disorders of the
5 eyelids; disorders of the conjunctiva; disorders of the
cornea; cataract; disorders of the uveal tract; disorders
of the retina; disorders of the optic nerve or visual
pathways; free radical induced eye disorders and
diseases; immunologically-mediated eye disorders and
10 disorders; eye injuries; and symptoms and complications
of eye disease, eye disorder, or eye injury.

17. The pharmaceutical composition of claim 12,
which is for improving naturally-occurring vision in an
15 animal, in the absence of any ophthalmologic disorder,
disease, or injury.

18. The pharmaceutical composition of claim 12,
wherein the N-heterocyclic carboxylic acid or carboxylic
20 acid isostere comprises a compound of formula (I):



I

where

n is 1-3;

X is either O or S;

R₁ is selected from the group consisting of C₁-C₉,
5 straight or branched chain alkyl or alkenyl, C₂-C₉,
straight or branched chain alkenyl, aryl, heteroaryl,
carbocycle, or heterocycle;

D is a bond, or a C₁-C₁₀ straight or branched chain
alkyl, C₂-C₁₀ alkenyl or C₂-C₁₀ alkynyl; and

10 R₂ is carboxylic acid or a carboxylic acid isostere;
wherein said alkyl, alkenyl, alkynyl, aryl, heteroaryl,
carbocycle, or heterocycle is optionally substituted with
one or more substituents selected from R³, where

R³ is hydrogen, hydroxy, halo, haloalkyl, thiocarbonyl,
15 alkoxy, alkenoxy, alkylaryloxy, aryloxy, arylalkyloxy,
cyano, nitro, imino, alkylamino, aminoalkyl, sulfhydryl,
thioalkyl, alkylthio, sulfonyl, C₁-C₆ straight or branched
chain alkyl, C₂-C₆ straight or branched chain alkenyl or
alkynyl, aryl, heteroaryl, carbocycle, heterocycle, and
20 CO₂R⁴ where R⁴ is hydrogen or C₁-C₉ straight or branched
chain alkyl or alkenyl;


or a pharmaceutically acceptable salt, ester, or solvate
thereof.

25 19. The pharmaceutical composition of claim 18,
wherein R₂ is a carbocycle or heterocycle containing any

combination of CH_2 , O, S, or N in any chemically stable oxidation state, wherein any of the atoms of said ring structure are optionally substituted in one or more positions with R^3 .

5

20. The pharmaceutical composition of claim 18, wherein R_2 is selected from the following group:



where the atoms of said ring structure may be optionally substituted at one or more positions with R^3 .

21. The pharmaceutical composition of claim 18,
5 wherein R_2 is selected from the group consisting of:
-COOH; -SO₃H,; -SO₂HNR³; -PO₂(R³)₂; -CN; -PO₃(R³)₂; -OR³; -
SR³; -NHCOR³; -N(R³)₂; -CON(R³)₂; -CONH(O)R³; -CONHNHSO₂R³;
-COHNSO₂R³; and -CONR³CN.


10 22. The pharmaceutical composition of claim 18,
wherein the N-heterocyclic carboxylic acid or carboxylic
acid isostere compound is selected from the group
consisting of compounds 1-138.

ABSTRACT OF THE DISCLOSURE


This invention relates to novel compositions and
uses of N-heterocyclic carboxylic acids and carboxylic
5 acid isosteres for treating a vision disorder, improving
vision, treating memory impairment or enhancing memory
performance in an animal.

Figure 1


GPI 1046 protects ganglion cells against degeneration due to 1 hour of retinal ischemia
Fluorogold labelled retinal ganglion cells in wholemount, 28 days after ischemic episode



A. Labelled retinal ganglion cells in the
Normal central retina




B. 1 hour of retinal ischemia produces
extensive loss of ganglion cells




B. Administration of GPI 1046 1 hour before retinal ischemia
and for 4 days subsequently produces significant protection of vulnerable retinal ganglion cells


Figure 2
GPI 1046 Protects retinal ganglion cell axons and prevents myelin degeneration
in the optic nerve induced by 1 hour of complete retinal ischemia,
toluidine blue stained optic nerve cross sections, 630X



A. Normal optic nerve




B. Vehicle treated optic nerve 28 days
after 1 hour complete retinal ischemia



C. GPI 1046 treated optic nerve 28 days
after 1 hour complete retinal ischemia

Figure 3

**GPI 1046 administration for 28 days
provides only moderate protection of
axotomized retinal ganglion cells**



Fluorogold labelled RGCs 90 days following transection,
Treatment with vehicle alone for 1st 28 days

Fluorogold labelled RGCs 90 days following transection,
Treatment with GPI 1046 for 1st 28 days
Treatment with vehicle alone for 1st 28 days




Figure 4


**GPI 1046 prevents axonal degeneration in the proximal stump of the optic nerve
RT97 neurofilament immunohistochemistry,
optic nerve cross sections, 90 days after complete transection**




A. Sham



B. Optic nerve transection (ONT) 90 days survival



**C. optic nerve 90 days after transection,
GPI 1046 treatment days 1-28**




**D. optic nerve 90 days after transection,
GPI 1046 treatment days 1-14**


Figure 5

**GPI 1046 administration for 28 days provides only moderate protection
of axotomized retinal ganglion cells**

Fluorogold labelled retinal ganglion cells 90 days following transection




vehicle administered for 1st 28 days




GPI 1046 administered for 1st 28 days

**GPI 1046 administration for 28 days preserves optic nerve axons
of surviving retinal ganglion cells**

RT 97 neurofilament immunohistochemistry 90 days after transection




vehicle administered for 1st 28 days



GPI 1046 administered for 1st 28 days

Figure 6

**Preservation of myelin in the proximal stump of the optic nerve 90 days after transection
14 vs 28 days treatment with GPI 1046 10mg/kg s.c.**




Normal(sham) Optic nerve



90 days after optic nerve transection, vehicle treated



90 days after optic nerve transection, 14 days GPI 1046



90 days after optic nerve transection, 28 days GPI 1046

myelin basic protein immunohistochemistry (SMI-94), 20X






Figure 7
FKBP-12 immunohistochemistry labels oligodendroglia and axons in the normal optic nerve

Figure 8


**GPI 1046 treatment prevents myelin degeneration
in the distal stump of the optic nerve
Myelin basic protein immunohistochemistry 90 days after transection**




A. Normal optic nerve



**B. Distal optic nerve stump
90 days after complete transection**



**C. Distal optic nerve stump
90 days after complete transection
GPI 1046 administered 1-14 days
after transection**



**D. Distal optic nerve stump
90 days after complete transection
GPI 1046 administered 1-28 days
after transection**

Figure 9


**GPI 1046 decreases neovascularization and prevents neuronal loss
in the inner retinal in the Streptozotocin model of diabetic retinopathy**

**A. Normal
retina
Cross section
Cresyl violet**

**Outer Nuclear
layer (ONL)**

**Inner Nuclear
layer(INL)**

**Ganglion cell
layer (GCL)**




**B. retina from
Streptozotocin
/vehicle case**

ONL

INL

GCL




**C. Retina from
Streptozotocin
/GPI 1046 case**

ONL

INL

GCL



DECLARATION FOR PATENT APPLICATION

Atty Docket 22762XY4S

As a below-named inventor(s), I/we hereby declare that:

My/Our residence(s), post office address(es) and citizenship(s) is/are as stated below next to my/our name(s).

We believe we are the original first and joint inventors of the subject matter which is claimed, and for which a patent is sought on the invention entitled:

the specification of which: (check one)

☒ is attached hereto.

☐ was filed on _____, as Serial No. _____,

and was amended on _____ 19 _____ (if applicable).

I/We hereby state that I/we have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I/We acknowledge the duty to disclose information which is material to the patentability of this application as defined by 37 CFR § 1.56.

I/We hereby claim foreign priority benefits under 35 U.S.C. § 119 of any foreign application(s) for patent or inventor's certificate listed below, and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Applications:

(Application No.)	(Country)	(Day/Month/Year Filed)	Priority Claimed	
			<input type="checkbox"/> Yes	<input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No
(Application No.)	(Country)	(Day/Month/Year Filed)	<input type="checkbox"/> Yes	<input type="checkbox"/> No

I/We hereby appoint Gary M. Nath, Reg. No. 26,965; Irvin A. Lavine, Reg. No. 16,838; Karen Lee Orzechowski, Reg. No. 31,621; Harold L. Novick, Reg. No. 26,011; Suet M. Chong, Reg. No. 38,104; Todd E. Juneau, Reg. No. 40,669; Patricia M. Drost, Reg. No. 29,790; Lee C. Heiman, Reg. No. P-41,827; Donald L. Sandler, Reg. No. 19,237; and Robert G. Lev, Reg. No. 30,280, as my attorneys to prosecute this application and transact all business in the U.S. Patent and Trademark Office connected therewith.

Direct Telephone Calls to:

Gary M. Nath
(202) 775-8383

Send Correspondence to:
NATH & ASSOCIATES
Suite 750
1835 K Street, N.W.
Washington, D.C. 20006 U.S.A.

I/We hereby claim the benefit under 35 U.S.C. § 120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by 35 U.S.C. § 112, first paragraph, I/we acknowledge the duty to disclose material information as defined in 37 CFR § 1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(U.S. Application Serial No.)	(U.S. Filing Date)	(Status--patented, pending, abandoned)
(U.S. Application Serial No.)	(U.S. Filing Date)	(Status--patented, pending, abandoned)

I/We hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. § 1001 and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of sole or first inventor: Douglas T. Ross

Inventor's Signature X Douglas T. Ross

Date 8/13/98

Residence: 316 South Main Street, North Wales, Pennsylvania 19454

Citizenship: United States of America

Post Office Address: same as above

Full name of second inventor: Hansjörg Sauer

Inventor's Signature X Hansjörg Sauer

Date 8/13/98

Residence: 10617 Lorain Avenue, Silver Spring, Maryland 20901

Citizenship: Germany

Post Office Address: same as above

Full name of third inventor: Gregory S. Hamilton

Inventor's Signature X Gregory S. Hamilton

Date 8/13/98

Residence: 6501 Frederick Road, Catonsville, Maryland 21228

Citizenship: United States of America

Post Office Address: same as above

Full name of fourth inventor: Joseph P. Steiner

Inventor's Signature X Joseph P. Steiner

Date 8/13/98

Residence: 4150 Louisville Road, Finksburg, Maryland 21048

Citizenship: United States of America

Post Office Address: _____

Full name of fifth inventor: _____

Inventor's Signature _____

Date _____

Residence: _____

Citizenship: _____

Post Office Address: _____

Full name of sixth inventor: _____

Inventor's Signature _____

Date _____

Residence: _____

Citizenship: _____

Post Office Address: _____